

A new characterization and a recognition algorithm of Lucas cubes

Andrej Taranenko[†]

Faculty of Natural Sciences and Mathematics, University of Maribor, Maribor, Slovenia

received 16th June 2012, revised 3rd Sep. 2013, accepted 25th Sep. 2013.

Fibonacci and Lucas cubes are induced subgraphs of hypercubes obtained by excluding certain binary strings from the vertex set. They appear as models for interconnection networks, as well as in chemistry. We derive a characterization of Lucas cubes that is based on a peripheral expansion of a unique convex subgraph of an appropriate Fibonacci cube. This serves as the foundation for a recognition algorithm of Lucas cubes that runs in linear time.

Keywords: Lucas cubes, characterization, recognition algorithm

1 Introduction

Graphs based on binary strings are used as models for interconnection networks. Hypercube, being a popular interconnection scheme for multicomputers, has often been replaced by other models with similar properties comparable to those of hypercubes, where the number of vertices and edges in these alternative models does not increase as rapidly.

As such a model Fibonacci cubes have been defined in Hsu (1993); Hsu et al. (1993), followed by extended Fibonacci cubes (Wu, 1997) and Lucas cubes (Dedó et al., 2002). Various studies have been made on the structure and different properties of Fibonacci cubes, Lucas cubes as well as Fibonacci-like cubes, see (Klavžar, 2005; Klavžar et al., 2011; Klavžar and Peterin, 2007; Munarini et al., 2001; Taranenko and Vesel, 2007), to name a few. We also refer to the extensive survey on Fibonacci cubes by Klavžar (2013). Both, Fibonacci and Lucas cubes, also appear in connection with resonance graphs in chemistry (Taranenko and Vesel, 2007; Žigert and Berlič, 2012).

A *Fibonacci string* of length n is a binary string $b_0b_1 \dots b_{n-1}$ such that $b_i \cdot b_{i+1} = 0$, for $i = 0, 1, \dots, n-2$. Equivalently, it is a binary string of length n without two consecutive ones.

The *Fibonacci cube* Γ_n of order n has the Fibonacci strings as vertices, with two vertices being adjacent whenever they differ in exactly one coordinate. We also set $\Gamma_0 = K_1$. See Fig. 1 for Fibonacci cubes of order n , for $n = 0, 1, \dots, 5$, with appropriate Fibonacci strings assigned to the vertices (these are omitted for Γ_5 for the clarity of the figure).

[†]Email: andrej.taranenko@uni-mb.si




Fig. 1: Fibonacci cubes Γ_n , for $n = 0, 1, \dots, 5$.

A *Lucas string* of length n is a binary string $b_0b_1 \dots b_{n-1}$ such that $b_i \cdot b_{i+1} = 0$, for $i = 0, 1, \dots, n-1$, where indices are computed modulo n . In other words, Lucas string is a binary string without two consecutive ones and without one both in the first and last position.

The *Lucas cube* Λ_n of order n is the graph with Lucas strings as vertices, again, with two vertices being adjacent whenever they differ in exactly one coordinate. We set $\Lambda_0 = K_1$. See Fig. 2 for Lucas cubes of order n , for $n = 0, 1, \dots, 5$, with appropriate Lucas strings assigned to the vertices (again, these are omitted for Λ_5 for the clarity of the figure). Note, that vertices of Λ_n can be obtained from the vertices of Fibonacci cubes Γ_{n-1} and Γ_{n-3} as follows: $V(\Lambda_n) = 0V(\Gamma_{n-1}) \cup 10V(\Gamma_{n-3})0$ (Munarini et al., 2001).

It is known that Fibonacci cubes and Lucas cubes are median graphs (Klavžar, 2005). For median graphs several efficient recognition algorithms are known. To mention the first polynomial algorithm with complexity $O(nm)$ presented in Jha and Slutzki (1992) and the much more advanced $O((m \log n)^{1.41})$ algorithm stated in Hammack et al. (2011). This raises the natural question whether a faster recognition algorithm for special families of median graphs exists. For Fibonacci cubes this question was partially answered by presenting a recognition algorithm with the complexity $O(m \log n)$ (Taranenko and Vesel, 2007). Even more, Vesel (2013) recently answered the problem of existence of a linear recognition algorithm for Fibonacci cubes, presented in Klavžar (2013). For Lucas cubes no recognition algorithm exists to our knowledge and is presented in this paper.

The paper is structured as follows. The next section contains some basic definitions and results concerning median graphs and Lucas cubes. In Section 3 several structural properties and a characterization of Lucas cubes are given. Using these results in the final Section 4 a recognition algorithm for Lucas cubes is presented. The presented algorithm correctly recognizes a Lucas cube in linear time.

2 Preliminaries

The *hypercube* Q_n of order n is the graph with the vertex set containing all binary strings of length n , where two vertices are adjacent whenever the two strings differ in exactly one position. Isometric graphs




Fig. 2: Lucas cubes Λ_n , for $n = 0, 1, \dots, 5$.

of hypercubes are called *partial cubes*.

The *Fibonacci numbers* form a sequence of positive integers F_n , where $F_0 = 0$, $F_1 = 1$ and for $n \geq 2$ satisfy the recurrence $F_n = F_{n-1} + F_{n-2}$.

Since a Lucas string is a binary string without two consecutive ones and without one both in the first and last position, we can say that a Lucas string contains no two consecutive ones in a circular manner. Moreover, in the rest of the paper, when dealing with Lucas strings, we will always compute indices modulo n , even when not explicitly stated.

For a triple of vertices u, v and w of a given graph G , a vertex x of G is a *median* of u, v and w if x lies simultaneously on shortest paths joining u and v , u and w , and v and w , respectively. If G is connected and every triple of vertices admits a unique median, then G is a *median graph*. It is well known that median graphs are partial cubes (cf. (Hammack et al., 2011)).

Let G be a connected graph with $e = xy$ and $f = uv$ two edges in G . We say that e is in relation Θ to f if $d(x, u) + d(y, v) \neq d(x, v) + d(y, u)$. Θ is reflexive and symmetric, but need not be transitive. We denote its transitive closure by Θ^* . It was proved in (Winkler, 1984) that G is a partial cube if and only if G is bipartite and $\Theta = \Theta^*$.

For $X \subseteq V(G)$ we denote the subgraph of G induced by the set X with $G[X]$.

A subgraph H of a graph G is called *convex*, if it is connected and if any shortest path of G between two vertices of H is completely in H .

Let H be a fixed subgraph of a graph G , $H \subseteq G$. The *peripheral expansion* $pe(G; H)$ of G with respect to H is the graph obtained from the disjoint union of G and an isomorphic copy of H , in which every vertex of the copy of H is joined by an edge with the corresponding vertex of H .

For an edge ab of graph G we define:

- $W_{ab} = \{w \in V(G) \mid d(a, w) < d(b, w)\}$
- $W_{ba} = \{w \in V(G) \mid d(b, w) < d(a, w)\}$

- $F_{ab} = \{xy \in E(G) \mid x \in W_{ab} \text{ and } y \in W_{ba}\}$
- $U_{ab} = \{w \in W_{ab} \mid w \text{ is the end vertex of an edge in } F_{ab}\}$
- $U_{ba} = \{w \in W_{ba} \mid w \text{ is the end vertex of an edge in } F_{ab}\}$

Let ab be an edge of a median graph G for which $U_{ab} = W_{ab}$. Then $G[W_{ab}]$ is called a *peripheral subgraph* of G . A Θ -class E of a median graph G is called *peripheral*, if at least one of $G[W_{ab}]$ and $G[W_{ba}]$ is peripheral for $ab \in E$. E is *internal* if it is not peripheral.

We will need the following well known lemma, cf. (Hammack et al., 2011).

Lemma 2.1 *Let $e = ab$ be an edge of a connected bipartite graph G . Then*

- (i) $F_{ab} = \{f \mid f \in E(G), e\Theta f\}$,
- (ii) $G \setminus F_{ab}$ has exactly two components.

The next theorem characterizes median graphs.

Theorem 2.2 (Mulder, 1978) *Let ab be an edge of a connected, bipartite graph G . Then G is a median graph if and only if the following three conditions are satisfied:*

- (i) $G[U_{ab}]$ is convex in $G[W_{ab}]$ and $G[U_{ba}]$ in $G[W_{ba}]$.
- (ii) F_{ab} is a matching defining an isomorphism between $G[U_{ab}]$ and $G[U_{ba}]$.
- (iii) $G[W_{ab}]$ and $G[W_{ba}]$ are median graphs.

3 Characterization

Before we present the theorem that characterizes Lucas cubes, we will state some properties concerning this family of graphs.

Proposition 3.1 (Klavžar and Peterin, 2007) *Let E be a Θ -class of the Lucas cube Λ_n . Then $|E| = F_{n-1}$.*

Let 0^p denote a binary string of length $p \geq 0$ with all bits equal to zero, similarly 1^p is a binary string of length p with all bits equal to 1. Let x and y be two arbitrary binary strings, we write xy for the concatenation of x and y .

Obviously the vertex 0^n , has exactly n neighbours in Λ_n , moreover it is the only vertex of degree n in Λ_n (Klavžar et al., 2011).

Proposition 3.2 *All neighbours of the vertex 0^n in the Lucas cube Λ_n of order $n \geq 3$ are of degree $n - 2$.*

Proof: Let $v = 0^n$. All the neighbours of v are binary strings with exactly one 1. Consider such a vertex u with one 1, say at p^{th} position, $0 \leq p \leq n - 1$. Since the bits at positions $p - 1$ and $p + 1$ must be 0, there are $n - 3$ vertices with exactly two bits equal to 1, that are adjacent to u . The vertex v is also adjacent to u , therefore u is of degree $n - 2$. Since u was arbitrarily chosen, this concludes the proof. \square

While it is known (Munarini et al., 2001) that a Lucas cubes of order n is composed from two Fibonacci cubes of order $n - 1$ and $n - 3$ with some additional edges between the two, the next proposition characterizes, how these two Fibonacci cubes are induced.

Proposition 3.3 *Let $a = 0^n \in V(\Lambda_n)$ and $ab \in E(\Lambda_n)$. Then the following hold:*

- (i) $\Lambda_n[W_{ab}]$ is isomorphic to Γ_{n-1} .
- (ii) $\Lambda_n[W_{ba}]$ is isomorphic to Γ_{n-3} .

Proof: Since b has exactly one 1, assume it is at p^{th} position, $0 \leq p \leq n-1$. Then

$W_{ab} = \{x \in V(\Lambda_n) \mid \text{where } x \text{ has 0 at } p^{\text{th}} \text{ position}\}$ and

$W_{ba} = \{x \in V(\Lambda_n) \mid \text{where } x \text{ has 1 at } p^{\text{th}} \text{ position}\}$.

(i) Consider a vertex $x \in W_{ab}$. We know that the vertex x is of the form $b_0 \dots b_{p-1} 0 b_{p+1} \dots b_{n-1}$, which is a Lucas string. Then the binary string of length $n-1$ defined as $b_{p+1} \dots b_{n-1} b_0 \dots b_{p-1}$ is clearly a Fibonacci string, since binary strings $b_0 \dots b_{p-1}$ and $b_{p+1} \dots b_{n-1}$ contain no two consecutive 1s, and b_0 and b_{n-1} cannot both be 1, since they are at the first and the last position of a Lucas string. The set W_{ab} contains all Lucas string of the described form, therefore we obtain all Fibonacci strings of length $n-1$ in this manner. Since two vertices of W_{ab} are connected if they differ in exactly one position, and the bit at position p , in the original Lucas string, we removed with this construction was equal to 0 in all strings, W_{ab} induces a Fibonacci cube of order $n-1$ in Λ_n and the assertion follows.

(ii) Consider a vertex $x \in W_{ba}$. We know that the vertex x is of the form $b_0 \dots b_{p-1} 1 b_{p+1} \dots b_{n-1}$, which is a Lucas string. Moreover, $b_{p-1} = b_{p+1} = 0$. Again, we construct a binary string of the form $b_{p+2} \dots b_{n-1} b_0 \dots b_{p-2}$, which is of length $n-3$. The rest follows the same line of thought as in (i), therefore the assertion follows. \square

Proposition 3.4 *Let E be a Θ -class of the Lucas cube Λ_n . Then E is peripheral.*

Proof: Let $a = 0^n$ be the unique vertex of degree n in Λ_n . All the edges with one end-vertex equal to a belong to different Θ classes. Since there are n such edges and Λ_n contains n different Θ -classes, we can arbitrarily choose one such edge. Let b be a vertex with exactly one 1 at p^{th} position, $0 \leq p \leq n-1$. Let $e = ab \in E(\Lambda_n)$ and $E = \{f \in E(\Lambda_n) \mid e\Theta f\}$. We claim that E is peripheral with $W_{ba} = U_{ba}$. By Proposition 3.3 the graph $\Lambda_n[W_{ba}]$ is isomorphic to Γ_{n-3} . A vertex $x \in W_{ba}$ is adjacent (among others) to the vertex y which differs from x in exactly position p . Since W_{ba} contains all and only vertices with 1 at position p , all vertices in W_{ba} are adjacent to a vertex in W_{ab} . It follows by definition of U_{ba} that $U_{ba} = W_{ba}$ and therefore E is peripheral. \square

From Theorem 2.2 and Propositions 3.3 and 3.4 we immediately obtain the following corollary.

Corollary 3.5 *Let $a = 0^n \in V(\Lambda_n)$ and $ab \in E(\Lambda_n)$. Then the graph $\Lambda_n[U_{ab}]$ is isomorphic to Γ_{n-3} .*

Let H be a subgraph of a graph G . Then ∂H is the set of all edges xy of G with $x \in H$ and $y \notin H$.

Theorem 3.6 *There exist exactly one convex subgraph $H \subseteq \Gamma_n$ isomorphic to Γ_{n-2} , such that the peripheral expansion $\text{pe}(\Gamma_n; H)$ is isomorphic to the Lucas cube Λ_{n+1} .*

Proof: Let H be a convex subgraph of Γ_n such that $\text{pe}(\Gamma_n; H)$ is isomorphic to Λ_{n+1} (assuming such H exists).

Note that the peripheral expansion increases the degree of each vertex in the chosen subgraph H by 1 in the graph obtained by the peripheral expansion. Since Λ_{n+1} contains exactly one vertex of degree $n+1$, the vertex $0^n \in V(\Gamma_n)$ must be in H . Moreover, the vertices $10^{n-1} \in V(\Gamma_n)$ and $0^{n-1}1 \in V(\Gamma_n)$,

which are both of degree $n - 1$ cannot be in H , since by Proposition 3.2 all the neighbouring vertices of $a = 0^{n+1} \in V(\Lambda_{n+1})$ are of degree $n - 1$, these two would be of degree n in the graph $\text{pe}(\Gamma_n; H)$, they would of course both be adjacent to a in this graph.

Let $b = 10^{n-1} \in V(\Gamma_n)$. Suppose a vertex $x \in W_{ba}$ is in $V(H)$. Since H is connected and we have established that $a \in V(H)$, there exists a shortest path from x to a , which is completely in H (since H is convex). But $a \in W_{ab}$, therefore any shortest path from a to x must contain an edge uv from F_{ab} (due to Lemma 2.1). By the Convexity lemma (Imrich and Klavžar, 1998), H is convex if and only if no edge of ∂H is in relation Θ to any edge in H . The edge $ab \in \partial H$, moreover $ab \in F_{ab}$ and is therefore in relation Θ with $uv \in E(H)$, which is a contradiction. So none of the vertices in W_{ba} are in H .

By the same reasoning, with $c = 0^{n-1}1 \in V(\Gamma_n)$, none of the vertices from W_{ca} are in H .

Recall, that $W_{ba}(W_{ca})$ contains all the vertices with 1 in the first and the last position, respectively. Consider the set $S := V(\Gamma_n) \setminus (W_{ba} \cup W_{ca}) = W_{ab} \cap W_{ac}$. Since the graphs induced by W_{ab} and W_{ac} are convex in Γ_n , the intersection (the graph induced by S) is also convex in Γ_n . Then every vertex $s \in S$ is of the form $0b_1b_2 \dots b_{n-2}0$, moreover it clearly contains all such vertices where $b_1b_2 \dots b_{n-2}$ is a Fibonacci string. Since two vertices are adjacent if they differ in exactly one position it follows that S induces a Fibonacci cube of order $n - 2$. Set $H := \Gamma_n[S]$, which is clearly the unique subgraph induced on vertices of S isomorphic to Γ_{n-2} , so H exists.

It remains to prove, that $\text{pe}(\Gamma_n; H)$ is isomorphic to Λ_{n+1} . As seen, the vertices of H are all of the form $0b_1b_2 \dots b_{n-2}0$. Let all the vertices in the isomorphic copy of H (introduced by the peripheral expansion) have the same assigned Fibonacci string as the corresponding vertices in H . Add a 0 to the beginning all vertices of Γ_n , and a 1 to the beginning of all vertices in the isometric copy. In this way we obtain all Lucas strings of length $n + 1$. From the construction of the strings it is clear that two strings are connected in $\text{pe}(\Gamma_n; H)$ if and only if they differ in exactly one position. \square

The next theorem characterizes Lucas cubes.

Theorem 3.7 *Let G be a connected bipartite graph, $a \in V(G)$ of degree n and $ab \in E(G)$. G is isomorphic to Λ_n if and only if the following conditions are upheld:*

- (1) *All the neighbours of a are of degree $n - 2$.*
- (2) *$G[U_{ab}]$ is convex in $G[W_{ab}]$.*
- (3) *F_{ab} defines an isomorphism between $G[U_{ab}]$ and $G[U_{ba}]$.*
- (4) *$G[W_{ab}] (G[W_{ba}])$ is isomorphic to $\Gamma_{n-1} (\Gamma_{n-3})$.*
- (5) *$W_{ba} = U_{ba}$.*

Proof: Suppose G is isomorphic to Λ_n . Condition (1) is true by Proposition 3.2. Since Λ_n is a median graph (Klavžar, 2005), by Theorem 2.2 conditions (2) and (3) are also true. Proposition 3.3 settles the condition (4). From Proposition 3.4 and its proof the final condition is also satisfied.

For the reverse, suppose for a graph G all conditions are met. Let $a \in V(G)$ be the vertex of degree n , and $b \in V(G)$ be an arbitrary neighbour of a (which by assumption is of degree $n - 2$). Clearly, if no such a exists or if there exists a neighbour b of a , such that the degree of b differs from $n - 2$, G is not isomorphic to a Lucas cube of order n , therefore these conditions are necessary. Assign Fibonacci strings of length $n - 1$ to all vertices of $G[W_{ab}]$, we will refer to them as labels. From the convexity of $G[U_{ab}]$

in $G[W_{ab}]$ and the proof of Theorem 3.6 we can choose the labelling such that the vertices in $G[U_{ab}]$ are of the form $0b_1b_2 \dots b_{n-2}0$. The set F_{ab} defines an isomorphism between $G[U_{ab}]$ and $G[U_{ba}]$, so assign to the vertices in $G[U_{ba}]$ the same Fibonacci string as their corresponding vertex in $G[U_{ab}]$. Now, add a 0 and 1 at the left hand side to all labels in $G[W_{ab}]$ and $G[W_{ba}]$, respectively. Since $U_{ba} = W_{ba}$, all the vertices have been assigned a (different) label. Moreover, the construction of the labels implies, that they are all Lucas strings of length n . What is left to prove is that two vertices are adjacent whenever the assigned labels differ in exactly one position. This clearly holds for all pairs of vertices, when both are either in W_{ab} or W_{ba} . Suppose, $x \in U_{ab}$ and $y \in U_{ba}$. Then $xy \in E(G)$ whenever $xy \in F_{ab}$, it follows that the corresponding labels differ in exactly one position. By Lemma 2.1 the sets W_{ab} and W_{ba} are disjoint, therefore two vertices $x \in W_{ab} \setminus U_{ab}$ and $y \in U_{ba}$ are not adjacent in G . By construction, the labels of x and y differ in more than one position. It follows that G is isomorphic to Λ_n . This concludes our proof. \square

Note that condition (1) of Theorem 3.7 is necessary, since b has been arbitrarily chosen. There exists a graph $G = \text{pe}(\Gamma_n; H)$ such that H is a convex subgraph of Γ_n isomorphic to Γ_{n-2} where G is not isomorphic to Λ_{n+1} . See Fig. 3 for an example.




Fig. 3: A peripheral expansion $\text{pe}(\Gamma_4; H)$ where H is a convex subgraph isomorphic to Γ_2

4 Recognition algorithm

Theorem 3.7 is also good from algorithmic point of view since it serves as the basis for the algorithm presented with procedure LUCAS for recognition of Lucas cubes. Procedure $\text{Fibonacci}(G, n, uv)$ used in procedure LUCAS is from Vesel (2013) and returns ACCEPT if a given graph G is isomorphic to a Fibonacci cube of order n , and REJECT otherwise. The input parameter uv represents an edge of the input graph G with one end-vertex of degree n . Moreover, it runs in $O(|E(\Gamma_n)|)$ time.

Before calling procedure LUCAS some preprocessing of the input graph is required. It is well known that $|V(\Gamma_h)| = F_{h+2}$. A given graph G , with $n = |V(G)|$ and $m = |E(G)|$, is examined only if

$n = F_{h+1} + F_{h-1}$ for some $h \geq 1$ and it is bipartite, otherwise graph is rejected. The computed value of h is also an input parameter of procedure LUCAS. The input graph fulfilling these conditions is declared isomorphic to a Lucas cube of order h if procedure LUCAS terminates without encountering REJECT statement.

In the presented procedure we denote by $N_S(v)$ the set of all neighbours of a vertex v , where neighbours are from the set S .

Procedure LUCAS (G, h)

Result: ACCEPT if G is isomorphic to Λ_h , REJECT otherwise

```

1 begin
2   if  $G$  is  $K_1$  or  $G$  is  $K_2$  then return ACCEPT
3   Find  $e = uv \in E(G)$  such that  $d(u) = h$  and  $d(v) = h - 2$ 
4   if  $e$  does not exist then return REJECT
5   Compute sets  $W_{uv}, W_{vu}, U_{uv}, U_{vu}$  and  $F_{uv}$ 
6   if  $F_{uv}$  is not a matching defining an isomorphism between  $G[U_{uv}]$  and  $G[U_{vu}]$  then
7     return REJECT
8   end
9   if  $G[U_{uv}]$  is not convex in  $G[W_{uv}]$  then return REJECT
10  if  $U_{vu} \neq W_{vu}$  then return REJECT
11   $u' \in N_{W_{uv}}(u)$ 
12   $v' \in N_{W_{vu}}(v)$ 
13  if Fibonacci( $G[W_{uv}], h - 1, uu'$ ) returns REJECT then return REJECT
14  if Fibonacci( $G[W_{vu}], h - 3, vv'$ ) returns REJECT then return REJECT
15  return ACCEPT
16 end
```

Since procedure LUCAS returns ACCEPT if and only if the conditions from Theorem 3.7 are satisfied, the following theorem follows immediately.

Theorem 4.1 Procedure LUCAS correctly recognizes a Lucas cube.

Theorem 4.2 Procedure LUCAS runs in $O(|m|)$ time to successfully recognize a Lucas cube Λ_h .

Proof: The steps from preprocessing can clearly be done in linear time. Note, that Steps 2-4 of procedure LUCAS can be computed in $O(m)$ time. It has been show in Jha and Slutzki (1992) that Steps 5-7 can be performed in time linear in the number of edges of the input graph. Moreover, Step 8 can be easily executed in linear time in the number of elements of the sets. Steps 9 and 10 can be done in constant time. So the complexity of the recognition algorithm depends on the complexity of the procedure Fibonacci(G, h, uv), which runs in linear time in the number of edges of the input graph (Vesel, 2013). So the time complexity of the procedure LUCAS is linear in the number of edges, ie. $O(|E(\Lambda_h)|)$. \square

Acknowledgements

This work has been financed by ARRS Slovenia under the grant P1-0297 and within the EUROCORES Programme EUROGIGA (project GReGAS) of the European Science Foundation. The author is also with Institute of Mathematics, Physics and Mechanics, Ljubljana.

References

- E. Dedó, D. Torri, and N. Z. Salvi. The observability of the fibonacci and the lucas cubes. *Discrete Math.*, 255:55–63, 2002.
- R. Hammack, W. Imrich, and S. Klavžar. *Handbook of Product Graphs, second edition*. CRC Press, Boca Raton, 2011.
- W. J. Hsu. Fibonacci cubes – a new interconnection topology. *IEEE Trans. Parallel Distr. Systems*, 4: 3–12, 1993.
- W. J. Hsu, C. V. Page, and J. S. Liu. Fibonacci cubes – a class of self-similar graphs. *Fibonacci Quart.*, 31:65–72, 1993.
- W. Imrich and S. Klavžar. A convexity lemma and expansion procedures for bipartite graphs. *European J. Combin.*, 19:677–685, 1998.
- P. K. Jha and G. Slutzki. Convex-expansions algorithms for recognition and isometric embeddings of median graphs. *Ars Combin.*, 34:75–92, 1992.
- S. Klavžar. On median nature and enumerative properties of fibonacci-like cubes. *Discrete Math.*, 299: 145–153, 2005.
- S. Klavžar. Structure of fibonacci cubes: a survey. *J. Comb. Optim.*, 25:505–522, 2013.
- S. Klavžar and I. Peterin. Edge-counting vectors, fibonacci cubes, and fibonacci triangle. *Publ. Math. Debrecen*, 71/3-4:267–278, 2007.
- S. Klavžar, M. Mollard, and M. Petkovšek. The degree sequence of fibonacci and lucas cubes. *Discrete Math.*, 311:1310–1322, 2011.
- H. M. Mulder. The structure of median graphs. *Discrete Math.*, 24:197–204, 1978.
- E. Munarini, C. P. Cippo, and N. Z. Salvi. On the lucas cubes. *Fibonacci Quart.*, 39:12–21, 2001.
- A. Taranenko and A. Vesel. Fast recognition of fibonacci cubes. *Algorithmica*, 49:81–93, 2007.
- A. Vesel. Linear recognition and embedding of fibonacci cubes. *Algorithmica*, to appear, 2013.
- P. M. Winkler. Isometric embeddings in products of complete graphs. *Discrete Appl. Math.*, 7:221–225, 1984.
- J. Wu. Extended fibonacci cubes. *IEEE Trans. Parallel Distr. Systems*, 8:3–9, 1997.
- P. Žigert and M. Berlič. Lucas cubes and resonance graphs of cyclic polyphenanthrenes. *MATCH Commun. Math. Comput. Chem.*, 68:79–90, 2012.

