Discrete Mathematics and Theoretical Computer Science DMTCS vol. **11:2**, 2009, 33–46

A construction of small regular bipartite graphs of girth 8

Camino Balbuena[1†]

[1]*Departament de Matemàtica Aplicada III, Universitat Politècnica de Catalunya, Campus Nord, Jordi Girona 1 i 3, 08034Barcelona, Spain*

received September 6, 2007, revised January 22, 2009, accepted June 25, 2009.

Let q be a prime a power and k an integer such that $3 \leq k \leq q$. In this paper we present a method using Latin squares to construct adjacency matrices of k-regular bipartite graphs of girth 8 on $2(kq^2 - q)$ vertices. Some of these graphs have the smallest number of vertices among the known regular graphs with girth 8.

Keywords: Incidence matrices, Latin squares, cages, girth.

1 Introduction

Throughout this paper, only undirected simple graphs without loops or multiple edges are considered. Unless otherwise stated, we follow the books by Godsil and Royle [16] and by Lint and Wilson [21] for terminology and definitions.

Let $G = (V(G), E(G))$ be a graph with vertex set $V = V(G)$ and edge set $E = E(G)$. The $girth$ of a graph G is the number $g = g(G)$ of edges in a smallest cycle. The *degree* of a vertex $v \in V$ is the number of vertices adjacent to v. A graph is called $regular$ if all the vertices have the same degree. A *cage* is a k-regular graph with girth g having the smallest possible number of vertices. Simply counting the numbers of vertices in the distance partition with respect to a vertex yields a lower bound $n_0(k, g)$ on the number of vertices $n(k, g)$ in a cage, with the precise form of the bound depending on whether g is even or odd.

$$n_0(k,g) = \begin{cases} 1 + k + k(k-1) + \ldots + k(k-1)^{(g-3)/2} & \text{if } g \text{ is odd;} \\ 2(1 + (k-1) + \ldots + (k-1)^{g/2-1}) & \text{if } g \text{ is even.} \end{cases} \tag{1}$$

As defined by Biggs [7], the *excess* of a k-regular graph G is the difference $|V(G)| - n_0(k, g)$. A (k, g)-cage with even girth g and $n_0(k, g)$ vertices is said to be a *generalized polygon graph*. Generalized polygon graphs exist if and only if $g \in \{4, 6, 8, 12\}$ [7]. The question of the construction of graphs with small excess is a difficult one. When $g = 6$, the existence of a graph with $n_0(k, 6) = 2(k^2 - k + 1)$

†This research was supported by the Ministry of Science and Innovation, Spain, and the European Regional Development Fund (ERDF) under project MTM2008-06620-C03-02/MTM.

1365–8050 © 2009 Discrete Mathematics and Theoretical Computer Science (DMTCS), Nancy, France

vertices called *generalized triangle*, is equivalent to the existence of a projective plane of order $k-1$, that is, a symmetric $((n_0(k,6))/2, k, 1)$-design. It is known that these designs exist whenever $k-1$ is a prime power, but the existence question for other values remains unsettled. *Generalized quadrangles* when $g = 8$, and *generalized hexagons* when $g = 12$ are also known to exist for all prime power values of $k-1$ [5, 16, 21].

Cages have been studied intensely since they were introduced by Tutte [27] in 1947. Erdős and Sachs [10] proved the existence of a graph for any value of the regularity k and the girth g, thus most of work carried out has been focused on constructing a smallest such graph [1, 2, 4, 6, 9, 11, 13, 15, 22, 23, 24, 28, 29]. Biggs is the author of an impressive report on distinct methods for constructing cubic cages [8]. For some time, Royle [26] kept a web-site in which all the cages known so far appear. More details about constructions on cages can be found in the survey by Wong [29], the survey by Holton and Sheehan [17] or on the more recent dynamic cage survey by Exoo and Jajcay [12].

It is conjectured that cages with even girth are bipartite [25, 29]. A graph is *bipartite* if its vertex set V can be partitioned into two partite sets, V_1 and V_2, such that any edge has one end in V_1 and the other in V_2. If the vertices are ordered in such a way that the vertices of V_1 come first, then the adjacency matrix of a bipartite graph can be written in the form

$$A = \begin{pmatrix} 0 & N \\ N^\top & 0 \end{pmatrix}. \tag{2}$$

An *incidence graph* is a bipartite graph in which the elements of one part V_1 are declared as *lines* and the elements of the other part V_2 are declared as *points*. The terminology for incidence graphs is geometric. A point and a line are said to be $incident$ if they are adjacent, thus the submatrix N of (2) is called an *incidence matrix* of the bipartite graph. If the number of points and the number of lines coincide, then N is clearly a square matrix. An incidence matrix N defines a *partial plane* when

- any line has at least two points, and
- two points are incident with at most one line.

Consequently, two lines of a partial plane have at most one point in common. The corresponding bipartite graph is called the *incidence graph of the partial plane*, which clearly has even girth $g \geq 6$. Thus for simplicity we shall say that *the partial plane has girth* $g/2$ if and only if the corresponding incidence graph has girth g.

Let q be a prime power and k an integer such that $3 \leq k \leq q$. In [3], incidence matrices of $(k,6)$-bipartite graphs of order $2(kq-1)$ were given. In this paper we present a method to construct the incidence matrices of k-regular bipartite graphs of girth $g = 8$ on $2q(kq-1)$ vertices.

2 Position matrices

Let S denote a set of symbols and let A be a matrix whose elements are subsets of S. Given $x \in S$ let $P_x(A)$ be a $(0,1)$-matrix of the same dimension as A that satisfies

$$(P_x(A))_{ij} = 1 \text{ if and only if } x \in A_{ij}.$$

Thus, $P_x(A)$ is called the *position matrix of the symbol* x in A. Suppose that $S = \{0, x_1, \ldots, x_n\}$. The position matrices of all the symbols in A different from 0 give rise to the following $(0,1)$-matrix $\mathcal{P}(A)$

called *position matrix of* A:

$$[\mathcal{P}(A)] = [P_{x_1}(A) \cdots P_{x_n}(A)].$$

Let $\{A^1, A^2, \ldots, A^r\}$ be a family of matrices of the same number of columns whose elements are subsets of S. Then the $(0,1)$-matrix spanned by the position matrices of all of them

$$\begin{pmatrix} \mathcal{P}(A^1) \\ \mathcal{P}(A^2) \\ \vdots \\ \mathcal{P}(A^r) \end{pmatrix} = \begin{pmatrix} P_{x_1}(A^1) & \cdots & P_{x_n}(A^1) \\ P_{x_1}(A^2) & \cdots & P_{x_n}(A^2) \\ & \vdots & \\ P_{x_1}(A^r) & \cdots & P_{x_n}(A^r) \end{pmatrix}, \tag{3}$$

is said to be *the position matrix of the family* $\mathcal{F} = \{A^1, A^2, \ldots, A^r\}$. The following example shows two matrices of order 2×2 whose elements are subsets of $S = \{0, a, b\}$ and the position matrix of them. From now on, if there is no confusion the 1–sets will be indicated as integers.

MATRICES			SYMBOLS			
			a		b	
A^1	a	a	**1**	**1**	0	0
	b	b	0	0	**1**	**1**
A^2	$\{a,b\}$	0	**1**	0	**1**	0
	0	$\{a,b\}$	0	**1**	0	**1**

(4)

As already mentioned in the Introduction, our main aim is to obtain incidence matrices of bipartite k-regular graphs of girth 8 with small excess. Such incidence matrices may be seen also as incidence matrices of partial planes which will be obtained by identifying row i of $P_z(A^\alpha)$ as *line* $i(\alpha)$, and column j of $P_z(A^\alpha)$ as *point* $j(z)$, for any matrix $A^\alpha \in \mathcal{F}$ and $z \in S - 0$. To achieve our goal we propose the following definitions.

Definition 2.1 *Let $g \geq 4$ be an even number. A family of matrices $\mathcal{F} = \{A^1, A^2, \ldots, A^r\}$ of the same number of columns whose elements are subsets of a set of symbols S is said to have girth g if the position matrix of $\mathcal{F}$ is the incidence matrix of a bipartite graph of girth g.*

Each matrix of (4) has girth $g = \infty$, and the two matrices A^1, A^2 form a family of girth 8.

Let us recall that a *Latin square* of order q is a $q \times q$ matrix with entries from a set of q symbols such that each symbol occurs exactly once in each row and exactly once in each column. A Latin square has clearly girth $g = \infty$ because the position matrices of its elements are permutation matrices yielding the incidence matrix of a partial plane consisting in a set of parallel lines (since they have no common point).

In [3] we introduced the notion of quasi row-disjoint matrices as follows.

Definition 2.2 *[3] Let A^1 and A^2 be two matrices of the same number of columns whose elements are subsets of a set of symbols S such that $0 \in S$. A pair (x, y) with $x, y \in S$ belongs to the cartesian product of any two rows $(A^1)_i \times (A^2)_h$ if and only if $(x, y) \in (A^1)_{ij} \times (A^2)_{hj}$ for some j. Then A^1 and A^2 are said to be* quasi row-disjoint *if and only if the cartesian product of any two rows $(A^1)_i, (A^2)_h$ contains at most one pair $(x, x) \in (A^1)_i \times (A^2)_h$ with $x \neq 0$.*

The pair of matrices of the example (4) are quasi row-disjoint matrices. Moreover, in [3] we have stated the following theorem which roughly speaking says that a family of r matrices is quasi row-disjoint if and only if its girth is at least 6.

Theorem 2.1 *[3] Let A^1 and A^2 be two matrices each one of girth at least 6 of the same number of columns and whose elements are subsets of a set of symbols S such that $0 \in S$. Then A^1 and A^2 are quasi row-disjoint if and only the family $\{A^1, A^2\}$ has girth at least 6.*

In the next theorem we give a sufficient condition for a family of matrices to have girth at least 8.

Theorem 2.2 *Let $\mathcal{F} = \{A^1, \ldots, A^r\}$ be a set of $r \geq 2$ quasi row-disjoint matrices of the same number of columns whose elements are subsets of a set of symbols S such that $0 \in S$. Let $(A^u)_i$, $(A^v)_{i'}$, $(A^w)_{i''}$ denote any three mutually distinct rows of matrices $A^u, A^v, A^w \in \mathcal{F}$. Then the girth of $\mathcal{F}$ is at least 8 if the sets $(A^u)_i \times (A^v)_{i'}, (A^v)_{i'} \times (A^w)_{i''}, (A^u)_i \times (A^w)_{i''}$ contains at most two distinct pairs (x, x), (y, y) with $x, y \neq 0$, $x, y \in S$.*

Proof: Suppose $(x, x) \in (A^u)_i \times (A^v)_{i'}$, $x \neq 0$. Therefore the position matrix of $\mathcal{F}$ has the following entries equal to 1:

$$P_x(A^u)(i, j) = P_x(A^v)(i', j) = 1, \tag{5}$$

where $P_x(A^u)$ and $P_x(A^v)$ are the position matrix of the element x in A^u and A^v respectively. Recall that for any given matrix $A^\alpha \in \mathcal{F}$ and $z \in S - 0$, row i of $P_z(A^\alpha)$ is line $i(\alpha)$, and column j of $P_z(A^\alpha)$ is point $j(z)$. Consequently, (5) means that lines $i(u)$ and $i'(v)$ have point $j(x)$ as a common point.

Analogously, $(y, y) \in (A^v)_{i'} \times (A^w)_{i''}$, $y \neq 0$, is equivalent to:

$$P_y(A^v)(i', j') = P_y(A^w)(i'', j') = 1,$$

or in other words, lines $i'(v)$ and $i''(w)$ have the point $j'(y)$ in common, with $j'(y) \neq j(x)$ because $y \neq x$. Thus if there exists $z \neq 0$, $z \neq x, y$, such that $(z, z) \in (A^u)_i \times (A^w)_{i''}$, then lines $i(u)$ and $i''(w)$ have the point $j''(z)$ in common, $j'' \neq j, j'$, yielding that the partial plane defined by the position matrix of $\mathcal{F}$ contains the triangle $j(x)j'(y)j''(z)$. In other words, the position matrix of $\mathcal{F}$ is the incidence matrix of a bipartite graph of girth less than 8. □

Our immediate goal is to derive a method for constructing a family of matrices with girth 8, because the position matrix of this family will be the incidence matrix of a bipartite graph of girth 8.

3 Method

Throughout this work let $[[n]]$ denote the set of non negative integers $\{0, 1, \ldots, n\}$ and $(n]] = [[n]] \setminus \{0\}$. Let I_n be the identity matrix and denote by $(t \times F)I_n$ the matrix obtained from I_n by replacing each one with a subset $\{t\} \times F$ of $\{t\} \times [[n]]$ for some $t \in [[n]]$ and $F \subseteq [[n]]$. In the following theorem we demonstrate a method for obtaining a family of matrices with girth 8 using Latin squares.

Theorem 3.1 *Let q be a prime power, and let $\mathbb{F}_q$ be the Galois field of order q. For each $u, t \in \mathbb{F}_q$, define the $q \times q$ matrix $L^{u,t}$ by*

$$L^{u,t}(i, j) = i + uj + ut, \;\; i, j \in \mathbb{F}_q.$$

Then the following assertions hold:

(i) For all $u, v \in \mathbb{F}_q \setminus \{0\}$, $u \neq v$, and $t, t' \in \mathbb{F}_q$ the matrices $L^{u,t}$ and $L^{v,t'}$ are quasi row-disjoint Latin squares with entries from $[[q-1]]$.

(ii) For any given $u, t \in \mathbb{F}_q$, $u \neq 0$, the matrix $L^{u,t} \times L^{u+u^{-1},t}$ has q^2 distinct entries. Moreover, the position matrix of the family

$$\{L^{u,t} \times L^{u+u^{-1},t} : t \in \mathbb{F}_q\}$$

is a $(0,1)$-matrix of order $q^2 \times q^3$ considering symbol $(0,0)$ different from 0, and it is the incidence matrix of a partial plane consisting in q^2 parallel lines with q points on each.

(iii) The family $\{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q, u \neq 0\}$ has girth 8 and its position matrix has order $(q^3 - q^2) \times q^3$ and q ones in each row and $q-1$ ones in each column.

(iv) The position matrix of the family

$$\{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q, u \neq 0\} \cup \{\, (t \times [[q-1]])I_q : t \in \mathbb{F}_q\}$$

is the incidence matrix of a q-regular bipartite graph of girth 8 with q^3 vertices in each partite set.

Proof: (i) Clearly $L^{u,t}$ and $L^{v,t'}$ are $q \times q$ Latin squares on $[[q-1]]$. Let us show that they are quasi row-disjoint. Otherwise, there exists $i, i' j, j' \in \mathbb{F}_q$ with $j \neq j'$ such that

$$\begin{aligned} L^{u,t}(i,j) &= L^{v,t'}(i',j), \\ L^{u,t}(i,j') &= L^{v,t'}(i',j'). \end{aligned}$$

Equivalently,

$$\begin{aligned} i + uj + ut &= i' + vj + vt', \\ i + uj' + ut &= i' + vj' + vt'. \end{aligned}$$

Therefore $u(j - j') = v(j - j')$ implying that $u = v$ or $j = j'$, a contradiction in either case.

(ii) Note that $L^{u+u^{-1},t}$ is also a Latin square if $u + u^{-1} \neq 0$, otherwise $L^{0,t}(i,j) = i$. In either case it is very easy to check that $L^{u,t}$ and $L^{u+u^{-1},t}$ are orthogonal, which implies that $L^{u,t} \times L^{u+u^{-1},t}$ has q^2 distinct entries. Hence, the position matrix of the family $\{L^{u,t} \times L^{u+u^{-1},t} : t \in \mathbb{F}_q\}$ is a $(0,1)$-matrix of order $q^2 \times q^3$ by considering entry (0,0) different from 0.

Let us show now that for every $i, i', j, t, t' \in \mathbb{F}_q$ such that $t \neq t'$ we have

$$(L^{u,t} \times L^{u+u^{-1},t})(i,j) \neq (L^{u,t'} \times L^{u+u^{-1},t'})(i',j).$$

Otherwise we would have

$$\begin{aligned} i + uj + ut &= i' + uj + ut', \\ i + (u+u^{-1})j + (u+u^{-1})t &= i' + (u+u^{-1})j + (u+u^{-1})t', \end{aligned}$$

implying $t = t'$, which is a contradiction. As a consequence, the position matrix of the set $\{L^{u,t} \times L^{u+u^{-1},t} : t \in \mathbb{F}_q\}$ has one unique entry equal to 1 in each column. Considering the rows of this $(0,1)$-matrix as lines and the columns as points, this is equivalent to say that each point belongs to a unique line. Thus, this $(0,1)$-matrix is the incidence matrix of q^2 parallel lines with q points on each.

(iii) First, let us show that the position matrix of the family $\mathcal{M} = \{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q,\ u \neq 0\}$ has q entries equal to 1 in each row and $q-1$ entries equal to 1 in each column. By (ii) the position matrix of each set $\{L^{u,t} \times L^{u+u^{-1},t} : t \in \mathbb{F}_q\}$ contributes with a unique 1 in each column, yielding that each column of the position matrix of $\mathcal{M}$ has $q-1$ entries equal to 1. Since each matrix $L^{u,t} \times L^{u+u^{-1},t}$ has q^2 distinct entries, then the q position matrices of symbols starting with the same x for any $x \in \mathbb{F}_q$ contribute with one unique 1 in each row. So the position matrix of $\mathcal{M}$ has q entries equal to 1 in each row. Therefore, we conclude that the position matrix of $\mathcal{M}$ has order $(q^3 - q^2) \times q^3$ and q entries equal to 1 in each row and $q-1$ entries equal to 1 in each column.

Next, let us show that the girth of $\mathcal{M}$ is at least 6. Otherwise, there exists $u, v \in \mathbb{F}_q \setminus \{0\}$, $u \neq v$ and $i, i'j, j' \in \mathbb{F}_q$ with $j \neq j'$ for which

$$\begin{array}{rcl} (L^{u,t} \times L^{u+u^{-1},t})(i,j) & = & (L^{v,t'} \times L^{v+v^{-1},t'})(i',j), \\ (L^{u,t} \times L^{u+u^{-1},t})(i,j') & = & (L^{v,t'} \times L^{v+v^{-1},t'})(i',j'). \end{array}$$

This implies that $L^{u,t}$ and $L^{v,t'}$ are not quasi row-disjoint, a contradiction with item (i). Thus the family $\{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q,\ u \neq 0\}$ has girth at least 6. Next let us show that the girth is 8 applying Theorem 2.2. By way of contradiction assume that for three elements u, v, w of $\mathbb{F}_q \setminus \{0\}$ there exist three pairwise distinct columns j, j', j'' for which

$$\begin{array}{rcl} (L^{u,t} \times L^{u+u^{-1},t})(i,j) & = & (L^{v,t'} \times L^{v+v^{-1},t'})(i',j) \\ (L^{v,t'} \times L^{v+v^{-1},t'})(i',j') & = & (L^{w,t''} \times L^{w+w^{-1},t''})(i',j') \\ (L^{w,t''} \times L^{w+w^{-1},t''})(i'',j'') & = & (L^{u,t} \times L^{u+u^{-1},t})(i,j''). \end{array} \tag{6}$$

Note that u, v, w must be three distinct elements because by (ii) each matrix $L^{u,t} \times L^{u+u^{-1},t}$ has q^2 distinct entries. Then from the equality between the first coordinates we have:

$$\begin{array}{rcl} (u-v)j & = & i' - i + vt' - ut; \\ (v-w)j' & = & i'' - i' + wt'' - vt'; \\ (w-u)j'' & = & i - i'' + ut - wt''. \end{array}$$

Hence

$$(v-u)j + (w-v)j' = (w-u)j''. \tag{7}$$

Moreover, from equalities between the second coordinates in (6) and taking into account (7) we obtain

$$(v^{-1} - u^{-1})j + (w^{-1} - v^{-1})j' = (w^{-1} - u^{-1})j''.$$

Multiplying this equality by uvw we get

$$w(u-v)j + u(v-w)j' = v(u-w)j''. \tag{8}$$

Multiplying (7) by w we also obtain

$$w(v-u)j + w(w-v)j' = w(w-u)j''. \tag{9}$$

Thus adding both equalities (8) and (9) we have

$$(u-w)(v-w)j' = (v-w)(u-w)j''.$$

Taking into account that u, v, w are mutually distinct we get that $j' = j''$ which is a contradiction. Therefore $\mathcal{M} = \{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q, u \neq 0\}$ has girth 8 as claimed.

(iv) By (iii) and applying Theorem 2.2, we only need to prove that for all $u, v \in \mathbb{F}_q \setminus \{0\}$ with $u \neq v$, any three matrices $L^{u,t} \times L^{u+u^{-1},t}$, $L^{v,t'} \times L^{v+v^{-1},t'}$ and $(t'' \times [[q-1]])I_q$ have girth 8. Otherwise we would have

$$\begin{array}{lcl} L^{u,t} \times L^{u+u^{-1},t}(i,j) & = & L^{v,t'} \times L^{v+v^{-1},t'}(i',j) \\ L^{v,t'} \times L^{v+v^{-1},t'}(i',j') & \in & (t'' \times [[q-1]])I_q(j',j') \\ L^{u,t} \times L^{u+u^{-1},t}(i,j') & \in & (t'' \times [[q-1]])I_q(j',j') \end{array}$$

Then $L^{u,t}(i,j) = L^{v,t'}(i',j)$ and $L^{u,t}(i,j') = L^{v,t'}(i',j') = t''$, meaning that $L^{u,t}$ and $L^{v,t'}$ are not quasi row-disjoint, contradicting item (i). Further, by (iii, we know that the position matrix of $\{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q, u \neq 0\}$ has q ones in each row and $q-1$ ones in each column. Since the rows of the position matrix of $(t \times [[q-1]])I_q$ contributes with one additional one, then the result follows. □

To illustrate the method of Theorem 3.1, both the matrices provided by this theorem and their position matrix for the first case $q = 2$ are shown in Table 1. From now on, if there is no confusion an entry (x, y) will be denoted as xy. Thus this (0,1)-matrix is the incidence matrix of a 2-regular graph of girth 8, which consists of two cycles of girth 8. In Table 2 the matrices for $q = 3$ are also depicted. The corresponding position matrix is the incidence matrix of a 3-regular graph of girth 8 on 27 vertices in each partite set. Table 3 contains the matrices for $q = 4$. Their position matrix is the incidence matrix of a 4-regular bipartite graph on 64 vertices in each partite set.

MATRICES			SYMBOLS			
			00	01	10	11
$L^{1,0} \times L^{0,0}$	00	10	**1** 0	0 0	0 **1**	0 0
	11	01	0 0	0 **1**	0 0	**1** 0
$L^{1,1} \times L^{0,1}$	10	00	0 **1**	0 0	**1** 0	0 0
	01	11	0 0	**1** 0	0 0	0 **1**
$(0 \times [[1]])I_2$	$\{00, 01\}$	0	**1** 0	**1** 0	0 0	0 0
	0	$\{00, 01\}$	0 **1**	0 **1**	0 0	0 0
$(1 \times [[1]])I_2$	$\{10, 11\}$	0	0 0	0 0	**1** 0	**1** 0
	0	$\{10, 11\}$	0 0	0 0	0 **1**	0 **1**

Tab. 1: Case $q = 2$.

Let us call *array of r symbols and n columns* the matrix of order $r \times n$

$$O_{r,n} = \begin{pmatrix} 1 & \cdots & 1 \\ 2 & \cdots & 2 \\ \vdots & \vdots & \vdots \\ r & \cdots & r \end{pmatrix}.$$

When $r = n$ the array is denoted by O_n. It is easy to see that the position matrix of $O_{r,n}$ is the incidence matrix of a partial plane consisting in r parallel lines, each one having n points. Using the position matrices of these arrays denoted by $\mathcal{P}(O_{r,n})$ and applying Theorem 3.1, we now present the method for constructing the desired $(k, 8)$-bipartite graphs.

	$L^{1,t} \times L^{2,t}$	$L^{2,t} \times L^{1,t}$	$(t \times [[2]])I_3$		
t=0	00 12 21	00 21 12	$\{00, 01, 02\}$	0	0
	11 20 02	11 02 20	0	$\{00, 01, 02\}$	0
	22 01 10	22 10 01	0	0	$\{00, 01, 02\}$
t=1	12 21 00	21 12 00	$\{10, 11, 12\}$	0	0
	20 02 11	02 20 11	0	$\{10, 11, 12\}$	
	01 10 22	10 01 22	0	0	$\{10, 11, 12\}$
t=2	21 00 12	12 00 21	$\{20, 21, 22\}$	0	0
	02 11 20	20 11 02	0	$\{20, 21, 22\}$	0
	10 22 01	01 22 10	0	0	$\{20, 21, 22\}$

Tab. 2: Matrices for the case $q = 3$.

	$L^{1,t} \times L^{0,t}$	$L^{2,t} \times L^{1,t}$	$L^{3,t} \times L^{1,t}$	$(t \times [[3]])I_4$			
t=0	00 10 20 30	00 21 32 13	00 31 12 23	$\{00, 01, 02, 03\}$	0	0	0
	11 01 31 21	11 30 23 02	11 20 03 32	0	$\{00, 01, 02, 03\}$	0	0
	22 32 02 12	22 03 10 31	22 13 30 01	0	0	$\{00, 01, 02, 03\}$	0
	33 23 13 03	33 12 01 20	33 02 21 10	0	0	0	$\{00, 01, 02, 03\}$
t=1	10 00 30 20	21 00 13 32	31 00 23 12	$\{10, 11, 12, 13\}$	0	0	0
	01 11 21 31	30 11 02 23	20 11 32 03	0	$\{10, 11, 12, 13\}$	0	0
	32 22 12 02	03 22 31 10	13 22 01 30	0	0	$\{10, 11, 12, 13\}$	0
	23 33 03 13	12 33 20 1	02 33 10 21	0	0	0	$\{10, 11, 12, 13\}$
t=2	20 30 00 10	32 13 00 21	12 23 00 31	$\{20, 21, 22, 23\}$	0	0	0
	31 21 11 01	23 02 11 30	03 32 11 20	0	$\{20, 21, 22, 23\}$	0	0
	02 12 22 32	10 31 22 03	30 01 22 13	0	0	$\{20, 21, 22, 23\}$	0
	13 03 33 23	01 20 33 12	21 10 33 02	0	0	0	$\{20, 21, 22, 23\}$
t=3	30 20 10 00	13 32 21 00	12 23 00 31	$\{30, 31, 32, 33\}$	0	0	0
	21 31 01 11	02 23 30 11	03 32 11 20	0	$\{30, 31, 32, 33\}$	0	0
	12 02 32 22	31 10 03 22	30 01 22 13	0	0	$\{30, 31, 32, 33\}$	0
	03 13 23 33	20 01 12 33	21 10 33 02	0	0	0	$\{30, 31, 32, 33\}$

Tab. 3: Matrices for the case $q = 4$.

Theorem 3.2 *Let q be a power prime, and let $\mathbb{F}_q$ be the Galois field of order q. For each $u, t \in \mathbb{F}_q$, $u \neq 0$, let $L^{u,t}$ be the matrix $L^{u,t}(i,j) = i + uj + ut,\ \ i, j \in \mathbb{F}_q$, and let M be the position matrix of the family*

$$\{L^{u,t} \times L^{u+u^{-1},t} : u, t \in \mathbb{F}_q, u \neq 0\} \cup \{\, (t \times [[q-1]])I_q : t \in \mathbb{F}_q\}.$$

Then the following assertions hold:

(i) The $(0,1)$-matrix

$$\boxed{M \;\big|\; \mathcal{P}(O_{q^2,q})^\top} \tag{10}$$

is the incidence matrix of a bipartite graph of girth 8 with $q^3 + q^2$ vertices in one partite set having degree $q + 1$, and q^3 vertices in the other partite set having degree q.

(ii) Let $(L^{u,t} \times L^{u+u^{-1},t})_0$ denote the matrix obtained from $L^{u,t} \times L^{u+u^{-1},t}$ by replacing each entry $0x$ with 0 for all $x \in \mathbb{F}_q$. Let M_0 be the position matrix of the family

$$\{(L^{u,t} \times L^{u+u^{-1},t})_0 : u, t \in \mathbb{F}_q, u \neq 0\} \cup \{\, (t \times [[q-1]])I_q : t \in \mathbb{F}_q, t \neq 0\}.$$

Then

$$\begin{array}{|c|c|}\hline M_0 & \mathcal{P}(O_{q^2-q,q})^\top \\ \hline & [0] \\ \hline \end{array} \tag{11}$$

is the incidence matrix of a q-regular bipartite graph of girth 8 with $q^3 - q$ vertices in each partite set.

(iii) Let k be an integer such that $3 \leq k \leq q-1$ and let $(L^{u,t} \times L^{u+u^{-1},t})_{q-k}$ denote the matrix obtained from $L^{u,t} \times L^{u+u^{-1},t}$ by replacing with 0 the entries $0y$ and $(x, x+s)$ for all $x, y, s \in \mathbb{F}_q$ for $s = 0, \ldots, q-1-k$. Let M_{q-k} be the position matrix of

$$\{(L^{u,t} \times L^{u+u^{-1},t})_{q-k} : u = 1, \ldots, k-1\} \cup \{\, (t \times ([[q-1]] \setminus \{t, t+1, \ldots, t+q-1-k\}))I_q : t \neq 0\}.$$

Suppose that the q columns of $O_{kq-q,q}$ are indexed by $j \in \mathbb{F}_q$. Let $O^*_{kq-q,q}$ be the matrix obtained from $O_{kq-q,q}$ by changing for 0 the entries $(i, -u^2 s)$ for all $u = 1, \ldots, k-1$, $i = (u-1)q+1, \ldots, uq$ and $s = 0, \ldots, q-1-k$. Then

$$\begin{array}{|c|c|}\hline M_{q-k} & \mathcal{P}(O^*_{kq-q,q})^\top \\ \hline & [0] \\ \hline \end{array} \tag{12}$$

is the incidence matrix of a k-regular bipartite graph of girth 8 with $kq^2 - q$ vertices in each partite set.

Proof: (i) From Theorem 3.1, it follows that M is the incidence matrix of a bipartite graph of girth 8 with q^3 lines and q^3 points. Thus we need to prove that the q^2 columns of $\mathcal{P}(O_{q^2,q})^\top$ can be added to M without decreasing the girth 8. It is readily seen that after adding these columns, the girth is at least 6, because by Theorem 3.1, for each $u \in \mathbb{F}_q \setminus \{0\}$ the set $\{L^{u,t} \times L^{u+u^{-1},t} : t \in \mathbb{F}_q, \}$ consists of q^2 parallel lines. Thus suppose that (10) contains as a sub-matrix the incidence matrix of a cycle of length 6. Then there exists $u, v \in \mathbb{F}_q \setminus \{0\}$ and $i, i', i'', j, j' \in \mathbb{F}_q$ with $u \neq v$, $i \neq i'$ and $j \neq j'$ such that

$$\begin{array}{lcl} (L^{u,t} \times L^{u+u^{-1},t})(i,j) & = & (L^{v,t'} \times L^{v+v^{-1},t'})(i'', j), \\ (L^{u,t} \times L^{u+u^{-1},t})(i',j') & = & (L^{v,t'} \times L^{v+v^{-1},t'})(i'', j'). \end{array}$$

From the equalities between their coordinates we obtain

$$\begin{array}{lcl} i - i' + u(j-j') & = & v(j-j'), \\ i - i' + (u+u^{-1})(j-j') & = & (v+v^{-1})(j-j'). \end{array}$$

Hence $u^{-1}(j-j') = v^{-1}(j-j')$, implying $u = v$ or $j = j'$, a contradiction in either case. Further if $(L^{u,t} \times L^{u+u^{-1},t})(i,j) \in (t' \times [[q-1]]I_q)(i'', j)$, and $(L^{u,t} \times L^{u+u^{-1},t})(i',j') \in (t' \times [[q-1]]I_q)(i'', j')$, then then $i'' = j = j'$ which is also a contradiction.

MATRICES			SYMBOLS																					
			10			11			12			20			21			22						
0	12	21	0	0	0	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0	0	**1**	0	0	0	0
11	20	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0	0	0	0
22	0	10	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	**1**	0	0	0
12	21	0	0	0	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0	0	0	0	0	**1**	0	0
20	0	11	0	0	0	0	0	**1**	0	0	0	**1**	0	0	0	0	0	0	0	0	0	**1**	0	0
0	10	22	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	**1**	0	0
21	0	12	0	0	0	0	0	0	0	0	**1**	0	0	0	**1**	0	0	0	0	0	0	0	**1**	0
0	11	20	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0	0	0	0	0	0	0	0	**1**	0
10	22	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	0	0	**1**
0	21	12	0	0	0	0	0	0	0	0	**1**	0	0	0	0	**1**	0	0	0	0	0	0	**1**	0
11	0	20	0	0	0	**1**	0	0	0	0	0	0	0	**1**	0	0	0	0	0	0	0	**1**	0	0
22	10	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	0	0	**1**	0
21	12	0	0	0	0	0	0	0	0	**1**	0	0	0	0	**1**	0	0	0	0	0	0	0	0	**1**
0	20	11	0	0	0	0	0	**1**	0	0	0	0	**1**	0	0	0	0	0	0	0	0	0	**1**	0
10	0	22	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	0	**1**	0
12	0	21	0	0	0	0	0	0	**1**	0	0	0	0	0	0	0	**1**	0	0	0	0	0	0	**1**
20	11	0	0	0	0	0	**1**	0	0	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	**1**
0	22	10	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	**1**
$\{10,11,12\}$	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	$\{10,11,12\}$	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	$\{10,11,12\}$	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0	0	0	0	0	0	0	0	0	0
$\{20,21,22\}$	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0	0	0
0	$\{20,21,22\}$	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0	0
0	0	$\{20,21,22\}$	0	0	0	0	0	0	0	0	0	0	0	**1**	0	0	**1**	0	0	**1**	0	0	0	0

Tab. 4: Case $q = 3$ for (ii) of Theorem 3.2. Incidence matrix of a (3,8)-graph on 48 vertices.

(ii) This item follows directly from the fact that M_0 is a sub-matrix of M obtained by deleting the first q^2 columns, which correspond to the position of the symbols starting by 0. Hence M_0 has $q^3 - q^2$ columns. Moreover, the total number of rows of M_0 is the number of matrices $L^{u,t} \times L^{u+u^{-1},t}$, (that is, $(q-1)q$) plus the number of matrices $(t \times ([[q-1]]))I_q$, $t \neq 0$, (a total of $q-1$) multiplied by q, that is, $q(q^2 - q + q - 1) = q^3 - q$. Thus M_0 is a matrix of order $(q^3 - q) \times (q^3 - q^2)$. Since $\mathcal{P}(O_{q^2-q,q})^\top$ contributes with $q^2 - q$ more columns, then (11) is a square matrix of order $q^3 - q$. Reasoning as in (iii) of Theorem 3.1, we obtain that (11) is the incidence matrix of a bipartite graph of girth 8, which has $q^3 - q$ columns and $q^3 - q$ rows both having q ones, so this item is valid. By way of example, both the matrices provided by this item (ii) and their position matrix for the case $q = 3$ are shown in Table 4. Thus this (0,1)-matrix is the incidence matrix of a 3-regular graph of girth 8 on 24 vertices in each partite set.

(iii) Note that (12) is a sub-matrix of (10), then it is the incidence matrix of a bipartite graph of girth 8. Moreover, M_{q-k} is obtained from M by deleting the first q^2 columns which corresponds to the position matrix of symbols starting by 0, and by deleting also $(q-k)(q-1)q$ columns corresponding to the symbols $(x+s, x+2s)$ for all $s = 0, 1, \ldots, q-1-k$ which have been changed for 0. Then the total number of columns of M_{q-k} is

$$q^3 - q^2 - (q-k)(q-1)q = kq(q-1).$$

The total number of rows of M_{q-k} is given by the number of matrices $(L^{u,t} \times L^{u+u^{-1},t})_{q-k}$, $u = 1, \ldots, k-1$, $t \in \mathbb{F}_q$, plus the number of matrices $(t \times ([[q-1]] \setminus \{t, t+1, \ldots, t+q-1-k\}))I_q : t \neq 0$, $t \neq 0$, (a total of $q-1$) multiplied by q, that is

$$(k-1)q^2 + (q-1)q = kq^2 - q.$$

Thus M_{q-k} is a matrix of order $(kq^2 - q) \times (kq^2 - kq)$. Since $\mathcal{P}(O^*_{kq-q,q})^\top$ contributes with $kq - q$ more columns, then (12) is a square matrix of order $kq^2 - q$.

To finish the proof of this item, we only need to show that (12) has k ones in each row and k ones in each column. To see this, let us show that given a fixed $s = 0, 1, \ldots, q - 1 - k$, the entries $(y, y + s)$ for all $y \in \mathbb{F}_q$, are in the same column of each matrix $L^{u,t} \times L^{u+u^{-1},t}$. Suppose $L^{u,t} \times L^{u+u^{-1},t}(i, j) = (y, y + s)$, that is

$$i + uj + ut = y, \text{ and } i + (u + u^{-1})j + (u + u^{-1})t = y + s.$$

Then $u^{-1}(j + t) = s$, which implies $j = us - t$. Thus our claim is true since the symbols $(y, y + s)$ are placed in the same column $us - t$ of the matrix $L^{u,t} \times L^{u+u^{-1},t}$. Therefore, after changing for 0 the symbols $0x$ and $(y, y+s)$ for all $x, y \in \mathbb{F}_q$ and for all $s = 0, \ldots, q-1-k$ we obtain a new matrix having k entries different from 0 in the rows i such that $i + u^2 s = 0$, and in the remaining rows $k - 1$ entries different from 0. Since in the rows i of $\mathcal{P}(O^*_{qk-q,q})^\top$ such that $i + u^2 s \neq 0$ there is a 1 by hypothesis, then $\mathcal{P}(O^*_{qk-q,q})^\top$ contributes with one additional entry equal to 1 in the same rows as those having $k-1$ entries different from 0. Hence (12) is a square matrix of order $kq^2 - q$ having k ones in each row and clearly in each column.

By way of example, both the matrices provided by this item (iii) and their position matrix for the case $q = 4$ and $k = 3$ are shown in Table 5; and the corresponding matrices for $q = 5$ and $k = 3$ are shown in Table 6. Thus, these (0,1)-matrices are the incidence matrix of a 3-regular graph of girth 8 on 44 vertices in each partite set for $q = 4$ and 70 vertices in each partite set for $q = 5$. □

4 Conclusion

For q a prime power and $3 \leq k \leq q$ we have presented a method providing the incidence matrices of k-regular bipartite graphs of girth 8 with $kq^2 - q$ vertices in each partite set. Thus if $n(k, 8)$ denotes the order of a $(k, 8)$-cage, it follows from (1) that

$$2k(k^2 - 2k + 2) \quad \leq \quad n(k, 8) \quad \leq \quad 2q(kq - 1).$$

Hence the q-regular bipartite graphs constructed in this work have an excess of $4q^2 - 6q$. And the $(q-1)$-regular bipartite graphs have an excess of $8q^2 - 20q + 10$.

As regards to known upper bounds on $n(k, g)$, Lazebnik, Ustimenko and Woldar [19] gave the following result: Let $k \geq 2$ and $g \geq 5$ be integers, and let q denote the smallest odd prime power for which $k \leq q$. Then,

$$n(k, g) \leq 2kq^{\frac{3}{4}g-a}\ , \tag{13}$$

where $a = 4, 11/4, 7/2, 13/4$ for $g \equiv 0, 1, 2, 3 \pmod 4$, respectively. According to (13), $n(k, 8) \leq 2kq^2$, therefore the graphs provided by our method also improve this result for $g = 8$. A construction giving this upper bound (13) for $g = 8$ appeared for the first time in [18] and was used later in [14] and probably, the simplest exposition of it is in Section 2.4 of [20]. In [15], $(q, 8)$-graphs with $2q(q^2 - 2)$ vertices were constructed using geometrical techniques. But for regularities $k < q$ the graphs constructed in this paper have the smallest number of vertices among the known regular graphs with girth 8.

				SYMBOLS									
				10	12	13	20	21	23	30	31	32	
0	10	20	30	0 1 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
0	0	31	21	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	1 0 0 0 0 0 0 0
0	32	0	12	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	1 0 0 0 0 0 0 0
0	23	13	0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0 0 0 0 0
10	0	30	20	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
0	0	21	31	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 1 0 0 0 0 0 0
32	0	12	0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 1 0 0 0 0 0 0
23	0	0	13	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0 0 0 0 0
20	30	0	10	0 0 0 1	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
31	21	0	0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 1 0 0 0 0 0
0	12	0	32	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 1 0 0 0 0 0
13	0	0	23	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0 0 0 0 0
30	20	10	0	0 0 1 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
21	31	0	0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 1 0 0 0 0
12	0	32	0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 1 0 0 0 0
0	13	23	0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1 0 0 0 0
0	21	32	13	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0 0 0 0 0
0	30	23	0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0 1 0 0 0
0	0	10	31	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0 1 0 0 0
0	12	0	20	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 1 0 0 0
21	0	13	32	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 0 0 0 0 0
30	0	0	23	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 1 0 0
0	0	31	10	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0 0 1 0 0
12	0	20	0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 1 0 0
32	13	0	21	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0 0 0 0 0
23	0	0	30	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0 0 0 1 0
10	31	0	0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0 0 0 1 0
0	20	0	12	0 0 0 0	0 0 0 1	0 0 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 1 0
13	32	21	0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 0 0 0 0 0 0
0	23	30	0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 1
31	10	0	0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0 0 0 0 1
20	0	12	0	0 0 0 0	0 0 1 0	0 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 1
				1 0 0 0	1 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 1 0 0	0 1 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
$\{10,\ 12,\ 13\}\ I_4$				0 0 1 0	0 0 1 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 1	0 0 0 1	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	1 0 0 0	1 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 1 0 0	0 1 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
$\{20,\ 21,\ 23\}\ I_4$				0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 1 0	0 0 1 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 1	0 0 0 1	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	1 0 0 0	1 0 0 0	1 0 0 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 1 0 0	0 1 0 0	0 1 0 0	0 0 0 0 0 0 0 0
$\{30,\ 31,\ 32\}\ I_4$				0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 1 0	0 0 1 0	0 0 1 0	0 0 0 0 0 0 0 0
				0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 0	0 0 0 1	0 0 0 1	0 0 0 1	0 0 0 0 0 0 0 0

Tab. 5: Case $q = 4$ for $k = 3$ of Theorem 3.2 (iii). Incidence matrix of a (3,8)-graph on 88 vertices.

	t=0	t=1	t=2	t=3	t=4
	0 0 24 31 43	0 24 31 43 0	24 31 43 0 0	31 43 0 0 24	43 0 0 24 31
	0 0 30 42 0	0 30 42 0 0	30 42 0 0 0	42 0 0 0 30	0 0 0 30 42
u=1	0 0 41 0 10	0 41 0 10 0	41 0 10 0 0	0 10 0 0 41	10 0 0 41 0
	0 0 0 14 21	0 0 14 21 0	0 14 21 0 0	14 21 0 0 0	21 0 0 0 14
	0 0 13 20 32	0 13 20 32 0	13 20 32 0 0	20 32 0 0 13	32 0 0 13 20
	0 20 0 10 30	20 0 10 30 0	0 10 30 0 20	10 30 0 20 0	30 0 20 0 10
u=2	0 31 0 21 41	31 0 21 41 0	0 21 41 0 31	21 41 0 31 0	41 0 31 0 21
	0 42 0 32 0	42 0 32 0 0	0 32 0 0 42	32 0 0 42 0	0 0 42 0 32
	0 0 0 43 13	0 0 43 13 0	0 43 13 0 0	43 13 0 0 0	13 0 0 0 43
	0 14 0 0 24	14 0 0 24 0	0 0 24 0 14	0 24 0 14 0	24 0 14 0 0
		$\{10,\ 13,\ 14\}\ I_5$	$\{20,\ 21,\ 24\}\ I_5$	$\{30,\ 31,\ 32\}\ I_5$	$\{41,\ 42,\ 43\}\ I_5$

$$O^*_{10,5} = \begin{array}{|ccccc|} \hline 0 & 1 & 1 & 1 & 0 \\ 0 & 2 & 2 & 2 & 0 \\ 0 & 3 & 3 & 3 & 0 \\ 0 & 4 & 4 & 4 & 0 \\ 0 & 5 & 5 & 5 & 0 \\ 0 & 0 & 6 & 6 & 6 \\ 0 & 0 & 7 & 7 & 7 \\ 0 & 0 & 8 & 8 & 8 \\ 0 & 0 & 9 & 9 & 9 \\ 0 & 0 & 10 & 10 & 10 \\ \hline \end{array}$$

Tab. 6: Matrices for $q = 5$ and $k = 3$ according to Case (iii) of Theorem 3.2.

Acknowledgements

I would like to thank the anonymous referees for their suggestions and for their valuable comments.

References

[1] M. ABREU, M. FUNK, D. LABBATE, V. NAPOLITANO, *On (minimal) regular graphs of girth 6*, Australasian Journal of Combinatorics 35 (2006), 119–132.

[2] G. ARAUJO, D. GONZÁLEZ MORENO, J.J. MONTELLANO, O. SERRA, *On upper bounds and conectivity of cages*, Australasian Journal of Combinatorics 38 (2007), 221–228.

[3] C. BALBUENA, *Incidence matrices of projective planes and other bipartite graphs of few vertices*, *Siam Journal of Discrete Maths.* 22(4) (2008), 1351–1363.

[4] E. BANNAI AND T. ITO, *On finite Moore graphs*, J. Fac. Sci. Univ. Tokyo, Sect. I A Math 20 (1973), 191–208.

[5] L.M. BATTEN, *Combinatorics of finite geometries*, Cambridge University Press, Cambridge, UK, 1997.

[6] C.T. BENSON, *Minimal regular graphs of girth eight and twelve*, Canad. J. Math 18 (1966), 1091–1094.

[7] N. BIGGS, *Algebraic Graph Theory*, Cambridge University Press, New York, 1996.

[8] N. BIGGS, *Construction for cubic graphs with large girth*, Electron. J. Combin. 5 (1998), #A1.

[9] G. BRINKMANN, B. D. MCKAY AND C. SAAGER, *The smallest cubic graphs of girth nine*, Combinatorics Probability and Computing 5, (1995), 1–13.

[10] P. ERDÖS AND H. SACHS, *Reguläre Graphen gegebener Taillenweite mit minimaler Knotenzahl*, Wiss. Z. Uni. Halle (Math. Nat.), 12 (1963), 251–257.

[11] G. EXOO, *A Simple Method for Constructing Small Cubic Graphs of Girths 14, 15 and 16*, Electron. J. Combin. 3 (1996).

[12] G. EXOO AND R. JAJCAY, *Dynamic Cage Survey*, Electron. J. Combin. 15 (2008) #DS16.

[13] W. FEIT AND G. HIGMAN, *The non-existence of certain generalized polygons*, J. Algebra 1 (1964), 114–131.

[14] Z. FUREDI, F. LAZEBNIK, A. SERESS, V.A. USTIMENKO AND J.A. WOLDAR, *Graphs of prescribed girth and bi-degree*, J. of Combin. Theory Series B 64(2) (1995), 228–239.

[15] A. GÁCS AND T. HÉGER, *On geometric constructions of (k, g)-graphs*, Contributions to Discrete Mathematics, 3(1) (2008) 63–80.

[16] C. GODSIL AND G. ROYLE, *Algebraic Graph Theory*, Springer, NY 2000.

[17] D.A. HOLTON AND J. SHEEHAN, *The Petersen Graph, Chapter 6: Cages*, Cambridge University (1993).

[18] F. LAZEBNIK, V.A. USTIMENKO, AND A.J. WOLDAR, *New constructions of bipartite graphs on m, n vertices with may edges and without small cycles*, J. of Combin. Theory Series B 61(1) (1994), 111–117.

[19] F. LAZEBNIK, V.A. USTIMENKO, AND A.J. WOLDAR, *New upper bounds on the order of cages*, Electron. J. Combin. 4 (1997), no. 2.

[20] F. LAZEBNIK, AND A.J. WOLDAR, *General properties of some families of graphs defined by systems of equations*, J. Graph Theory 38 (2001), 65–86.

[21] J. H. VAN LINT AND R. M. WILSON, *A course in Combinatorics*, Cambridge University Press, UK 1994.

[22] M. MERINGER, *Fast generation of regular graphs and construction of cages*, J. Graph Theory 30 (1999), 137–146.

[23] M. O'KEEFE AND P.K. WONG, *A smallest graph of girth 10 and valency 3*, J. of Combin. Theory Series B 29 (1980), 91–105.

[24] M. O'KEEFE AND P.K. WONG, The smallest graph of girth 6 and valency 7, J. Graph Theory 5 (1981), no. 1, 79–85.

[25] T. PISANSKI, M. BOBEN, D. MARUSIC, A. ORBANIC, A. GRAOVAC, *The* 10*-cages and derived configurations*, Discrete Mathematics 275 (2004), 265–276.

[26] G. ROYLE, http://www.cs.uwa.edu.au/∼gordon/data.html.

[27] W. T. TUTTE, *A family of cubical graphs*, Proc. Cambridge Philos. Soc. (1947), 459–474.

[28] P.K. WONG, *A regular graph of girth 6 and valency 11*, Internat. J. Math. Sci 9, (1986), 561–565.

[29] P. K. WONG, *Cages-a survey*, J. Graph Theory 6 (1982), 1–22.