

A polynomial realization of the Hopf algebra of uniform block permutations.

Rémi Maurice¹

¹*Institut Gaspard Monge, Université Paris-Est Marne-la-Vallée, 5 Boulevard Descartes, Champs-sur-Marne, 77454 Marne-la-Vallée cedex 2, France*

Abstract. We investigate the combinatorial Hopf algebra based on uniform block permutations and we realize this algebra in terms of noncommutative polynomials in infinitely many bi-letters.

Résumé. Nous étudions l'algèbre de Hopf combinatoire dont les bases sont indexées par les permutations de blocs uniformes et nous réalisons cette algèbre en termes de polynômes non-commutatifs en une infinité de bi-lettres.

Keywords: combinatorial Hopf algebras, uniform block permutations

1 Introduction.

For several years, Hopf algebras based on combinatorial objects have been thoroughly investigated. Examples include the Malvenuto-Reutenauer Hopf algebra **FQSym** whose bases are indexed by permutations [7, 4, 2], the Loday-Ronco Hopf algebra **PBT** whose bases are indexed by planar binary trees [6], the Hopf algebra of free symmetric functions **FSym** whose bases are indexed by standard Young tableaux [11].

The product and the coproduct, which define the structure of a combinatorial Hopf algebra, are, in general, rules of composition and decomposition given by combinatorial algorithms such as the shuffle and the deconcatenation of permutations, as in **FQSym** [4], of packed words, as in **WQSym** [9] or of parking functions, as in **PQSym** [10]. These can also be given by the disjoint union and the admissible cuts of rooted trees, as in the Connes-Kreimer Hopf algebra [3], or by concatenation and subgraphs/contractions of graphs [8].

Computing with these structures can be difficult, and polynomial realization can bring up important simplifications.

We can see a polynomial realization as the image by an injective Hopf algebra morphism from a combinatorial Hopf algebra to an algebra of polynomials (commutative or not).

The idea is to encode the combinatorial objects by polynomials, in such a way that the product of the combinatorial Hopf algebra become the ordinary product of polynomials, and that the coproduct be given by the disjoint union of alphabets, endowed with some extra structure such as an order relation. This can be done, *e.g.*, for **FQSym** [4], **WQSym** [9], **PQSym** [10] and Connes-Kreimer [5].

In this paper, we realize the Hopf algebra **UBP** of uniform block permutations introduced by Aguiar and Orellana [1]. We begin by recalling the preliminary notions of uniform block permutations in Section

2. We recall, in Section 3, the Hopf algebra structure of uniform block permutations, we translate this structure via the decomposition and we describe the dual structure. Using polynomial realizations of Hopf algebras based on permutations and set partitions, we realize this algebra in Section 4. We finish by finding the Hopf algebra **WQSym** as a quotient of **UBP** and the dual as a subalgebra of **UBP***.

Acknowledgements

The author would like to thank Jean-Yves Thibon for his advice and reading of this paper.

2 Definitions and decompositions.

Definition 1 A uniform block permutation of size n is a bijection between two set partitions of $\{1, \dots, n\}$ that maps each block to a block of the same cardinality.

Notation 1 Let $\mathcal{A} = \{A_1, A_2, \dots, A_k\}$ and $\mathcal{A}' = \{A'_1, A'_2, \dots, A'_k\}$ be two set partitions of the same size. We denote the uniform block permutation $f : \mathcal{A} \rightarrow \mathcal{A}'$ for which the image of A_i is A'_i by

$$\left(\begin{array}{c|c|c} A_1 & A_2 & \dots & A_k \\ \hline A'_1 & A'_2 & \dots & A'_k \end{array} \right).$$

Example 1 Here are all uniform block permutations of size 1 and 2:

$$\begin{aligned} n = 1 & \quad \left(\begin{array}{c} 1 \\ 1 \end{array} \right) \\ n = 2 & \quad \left(\begin{array}{c|c} 12 \\ \hline 12 \end{array} \right), \left(\begin{array}{c|c} 1 & 2 \\ \hline 1 & 2 \end{array} \right), \left(\begin{array}{c|c} 1 & 2 \\ \hline 2 & 1 \end{array} \right). \end{aligned}$$

In the case where the set partitions consist of singletons (i.e., $\mathcal{A} = \{\{1\}, \dots, \{n\}\}$), we find the permutations of size n .

Definition 2 Let f and g be two uniform block permutations. The concatenation of f and g , denoted by $f \times g$, is the uniform block permutation obtained by adding the size of f to all entries of g and concatenating the result to f .

Example 2

$$\left(\begin{array}{c|c} 13 & 2 \\ \hline 23 & 1 \end{array} \right) \times \left(\begin{array}{c|c|c} 13 & 2 & 46 & 5 \\ \hline 46 & 2 & 15 & 3 \end{array} \right) = \left(\begin{array}{c|c|c|c} 13 & 2 & 46 & 5 & 79 & 8 \\ \hline 23 & 1 & 79 & 5 & 48 & 6 \end{array} \right).$$

Definition 3 Let $f : \mathcal{A} \rightarrow \mathcal{A}'$ and $g : \mathcal{B} \rightarrow \mathcal{B}'$ be two uniform block permutations of the same size n . The composition $g \circ f : \mathcal{C} \rightarrow \mathcal{C}'$ of f and g is defined by the following process. The blocks C of the set partition \mathcal{C} are the subsets of $\{1, \dots, n\}$ which are minimal for the two properties

1. C is a union of blocks of \mathcal{A}
2. $f(C)$ is a union of blocks B_i of \mathcal{B} .

The image of C is the union of images $g(B_i)$.

Note that if f is a permutation, then the set partition \mathcal{C}' is \mathcal{B}' , and if g is a permutation, then the set partition \mathcal{C} is \mathcal{A} .

Example 3

$$\begin{aligned} \left(\begin{array}{c|c} 125 & 346 \\ \hline 236 & 145 \end{array} \right) \circ \left(\begin{array}{c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 \\ \hline 5 & 4 & 2 & 6 & 3 & 1 \end{array} \right) &= \left(\begin{array}{c|c} 136 & 245 \\ \hline 236 & 145 \end{array} \right), \\ \left(\begin{array}{c|c|c|c|c|c} 15 & 2 & 3 & 4 & 6 \\ \hline 46 & 2 & 1 & 5 & 3 \end{array} \right) \circ \left(\begin{array}{c|c|c|c} 14 & 2 & 36 & 5 \\ \hline 23 & 1 & 45 & 6 \end{array} \right) &= \left(\begin{array}{c|c|c} 14 & 236 & 5 \\ \hline 12 & 456 & 3 \end{array} \right). \end{aligned}$$

The data of a permutation and a set partition is enough to describe a uniform block permutation. Given a uniform block permutation $f : \mathcal{A} \rightarrow \mathcal{A}'$ of size n , there is a permutation σ in \mathfrak{S}_n such that

$$\mathcal{A}' = \bigcup_{H \in \mathcal{A}} \left\{ \bigcup_{x \in H} \{\sigma(x)\} \right\}.$$

Conversely, a set partition of the set $\{1, \dots, n\}$ and a permutation of \mathfrak{S}_n determine a uniform block permutation of size n . Any uniform block permutation can thus decompose (non-uniquely) as

$$f = \sigma \circ Id_{\mathcal{A}} = Id_{\mathcal{A}'} \circ \sigma. \quad (1)$$

The first equality expresses the fact that one can group the images σ_i whose indices are in the same block in \mathcal{A} , the second expresses the fact that one can group the images σ_i whose values are in the same block in \mathcal{A}' .

3 Algebraic structures.

In [1], Aguiar and Orellana introduced a Hopf algebra whose basis is the set of uniform block permutations. It will be denoted by **UBP**, standing for Uniform Block Permutations. Let us recall the convolution of two permutations. Let τ and τ' be two permutations:

$$\tau * \tau' = \sum_{\substack{\sigma = u \cdot v \\ \text{std}(u) = \tau \\ \text{std}(v) = \tau'}} \sigma$$

where the *standardization* is a process that associates a permutation with a word. The standardized of w is defined as the permutation having the same inversions as w . For example:

$$\text{std}(a_2 a_3 a_2 a_1 a_6 a_1 a_4) = 3541726,$$

$$12 * 21 = 1243 + 1342 + 1432 + 2341 + 2431 + 3421.$$

The product and the coproduct are defined as follows.

Definition 4 Let f and g be two uniform block permutations of respective sizes m and n . The convolution, denoted by $f * g$, is the sum of all uniform block permutations of the form

$$\left(\begin{array}{c|c|c|c} 1 & 2 & \dots & m+n \\ \hline \sigma_1 & \sigma_2 & \dots & \sigma_{m+n} \end{array} \right) \circ (f \times g).$$

where

$$\text{std}(\sigma_1 \dots \sigma_m) = 12 \dots m \text{ and } \text{std}(\sigma_{m+1} \dots \sigma_{m+n}) = 12 \dots n$$

(i.e., σ appears in $12 \dots m * 12 \dots n$).

Example 4

$$\begin{aligned} \begin{pmatrix} 1 \\ 1 \end{pmatrix} * \begin{pmatrix} 12 \\ 12 \end{pmatrix} &= \left[\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix} + \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix} + \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} \right] \circ \begin{pmatrix} 1 & 23 \\ 1 & 23 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 23 \\ 1 & 23 \end{pmatrix} + \begin{pmatrix} 1 & 23 \\ 2 & 13 \end{pmatrix} + \begin{pmatrix} 1 & 23 \\ 3 & 12 \end{pmatrix}. \end{aligned}$$

We denote by $\text{std}(\mathcal{E})$ the standardization of a collection of disjoint sets of integers obtained by numbering each integer a in \mathcal{E} by one plus the number of integers in \mathcal{E} smaller than a .

Example 5

$$\text{std}(\{\{1, 6\}, \{9, 13\}, \{3, 5, 12\}\}) = \{\{1, 4\}, \{5, 7\}, \{2, 3, 6\}\}.$$

We set

$$\mathcal{C}_{\{1, \dots, i\}} = \{H \in \mathcal{C}; H \subset \{1, 2, \dots, i\}\}, \mathcal{C}_{\{i+1, \dots, n\}} = \text{std}(\mathcal{C} - \mathcal{C}_{\{1, \dots, i\}})$$

and let $I(\mathcal{C})$ be the set of integers i such that

$$\{1, 2, \dots, n\} = \mathcal{C}_{\{1, \dots, i\}} \times \mathcal{C}_{\{i+1, \dots, n\}}$$

for any set partition \mathcal{C} .

Example 6 Let $\mathcal{C} = \{\{1, 3\}, \{2\}, \{4, 6, 7\}, \{5\}\}$ be a set partition:

$$\begin{aligned} \mathcal{C}_{\{1, \dots, 2\}} &= \{\{2\}\} & \mathcal{C}_{\{1, \dots, 3\}} &= \{\{1, 3\}, \{2\}\} \\ \mathcal{C}_{\{4, \dots, 7\}} &= \{\{1, 3, 4\}, \{2\}\} & I(\mathcal{C}) &= \{0, 3, 7\} \end{aligned}$$

Definition 5 Let $f : \mathcal{A} \rightarrow \mathcal{A}'$ be a uniform block permutation of size n . By Lemma (2.2) in [1], i is in $I(\mathcal{A}')$ if and only if there exists a unique permutation σ in \mathfrak{S}_n appearing in the convolution product $12 \cdots i * 12 \cdots n - i$ and unique uniform block permutations $f_{(i)}$ of size i and $f'_{(n-i)}$ of size $n - i$ such that

$$f = (f_{(i)} \times f'_{(n-i)}) \circ \left(\begin{array}{c|c|c|c} 1 & 2 & \cdots & n \\ (\sigma^{-1})_1 & (\sigma^{-1})_2 & \cdots & (\sigma^{-1})_n \end{array} \right).$$

The coproduct is then

$$\Delta(f) := \sum_{i \in I(\mathcal{A}')} f_{(i)} \otimes f'_{(n-i)}.$$

Example 7

$$\Delta \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) = () \otimes \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) + \left(\begin{array}{c|c} 13 & 245 \\ 15 & 234 \end{array} \right) \otimes \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) \otimes ()$$

since

$$\begin{aligned} \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) &= \left[() \times \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) \right] \circ \left(\begin{array}{c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array} \right) \\ &= \left[\left(\begin{array}{c|c} 13 & 245 \\ 15 & 234 \end{array} \right) \times \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right] \circ \left(\begin{array}{c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 6 & 4 & 5 \end{array} \right) \\ &= \left[\left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) \times () \right] \circ \left(\begin{array}{c|c|c|c|c|c} 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array} \right). \end{aligned}$$

One can rewrite this product and coproduct using the decomposition of a uniform block permutation. Given $f : \mathcal{A} \rightarrow \mathcal{A}'$ and $g : \mathcal{B} \rightarrow \mathcal{B}'$ two uniform block permutations, $f = \sigma \circ Id_{\mathcal{A}} = Id_{\mathcal{A}'} \circ \sigma$ and $g = \tau \circ Id_{\mathcal{B}} = Id_{\mathcal{B}'} \circ \tau$, we have

$$f * g = (\sigma * \tau) \circ Id_{\mathcal{A} \times \mathcal{B}} \quad (2)$$

$$\Delta(f) = \sum_{i \in I(\mathcal{A}')} f_{\{1, \dots, i\}} \otimes f_{\{i+1, \dots, n\}} \quad (3)$$

where

$$\begin{aligned} f_{\{1, \dots, i\}} &= Id_{\mathcal{A}'_{\{1, \dots, i\}}} \circ \sigma_{\{1, \dots, i\}} \\ f_{\{i+1, \dots, n\}} &= Id_{\mathcal{A}'_{\{i+1, \dots, n\}}} \circ \text{std}(\sigma_{\{i+1, \dots, n\}}). \end{aligned}$$

Example 8

$$\begin{aligned} \begin{pmatrix} 1 \\ 1 \end{pmatrix} * \begin{pmatrix} 12 \\ 12 \end{pmatrix} &= 1 \circ Id_{\{\{1\}\}} * 12 \circ Id_{\{\{1,2\}\}} \\ &= (1 * 12) \circ Id_{\{\{1\}, \{2,3\}\}} \\ &= (123 + 213 + 312) \circ Id_{\{\{1\}, \{2,3\}\}} \\ &= \begin{pmatrix} 1 & | & 23 \\ 1 & | & 23 \end{pmatrix} + \begin{pmatrix} 1 & | & 23 \\ 2 & | & 13 \end{pmatrix} + \begin{pmatrix} 1 & | & 23 \\ 3 & | & 12 \end{pmatrix}. \end{aligned}$$

Example 9 We set $\mathcal{A} = \{\{1, 5\}, \{2, 3, 4\}, \{6\}\}$.

$$\begin{aligned} \Delta \left(\begin{array}{c|cc} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) &= \Delta(Id_{\mathcal{A}} \circ 125634) \\ &= \varepsilon \otimes Id_{\mathcal{A}} \circ 125634 + Id_{\{\{1,5\}, \{2,3,4\}\}} \circ 12534 \otimes Id_{\{\{1\}\}} \circ 1 + Id_{\mathcal{A}} \circ 125634 \otimes \varepsilon \\ &= () \otimes \begin{pmatrix} 13 & | & 256 & | & 4 \\ 15 & | & 234 & | & 6 \end{pmatrix} + \begin{pmatrix} 13 & | & 245 \\ 15 & | & 234 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 13 & | & 256 & | & 4 \\ 15 & | & 234 & | & 6 \end{pmatrix} \otimes (). \end{aligned}$$

It is proved in [1] that **UBP** is a self-dual Hopf algebra. We can describe the dual structure. Let us first define another product and coproduct of uniform block permutations. Let $f : \mathcal{A} \rightarrow \mathcal{A}'$ and $g : \mathcal{B} \rightarrow \mathcal{B}'$ be two uniform block permutations, $f = \sigma \circ Id_{\mathcal{A}} = Id_{\mathcal{A}'} \circ \sigma$ and $g = \tau \circ Id_{\mathcal{B}} = Id_{\mathcal{B}'} \circ \tau$:

$$f \sqcap g = Id_{\mathcal{A}' \times \mathcal{B}'} \circ (\sigma \sqcap \tau) \quad (4)$$

$$\Delta_1(f) = \sum_{i \in I(\mathcal{A})} f_1 \cdots f_i \otimes f_{i+1} \cdots f_n \quad (5)$$

where

$$\begin{aligned} f_1 \cdots f_i &= \text{std}(\sigma_1 \cdots \sigma_i) \circ Id_{\mathcal{A}'_{\{1, \dots, i\}}} \\ f_{i+1} \cdots f_n &= \text{std}(\sigma_{i+1} \cdots \sigma_n) \circ Id_{\mathcal{A}'_{\{i+1, \dots, n\}}}. \end{aligned}$$

Example 10

$$\begin{aligned}
\begin{pmatrix} 1 \\ 1 \end{pmatrix} \bar{\sqcup} \begin{pmatrix} 12 \\ 12 \end{pmatrix} &= Id_{\{\{1\}, \{2,3\}\}} \circ (1 \bar{\sqcup} 12) \\
&= Id_{\{\{1\}, \{2,3\}\}} \circ (123 + 213 + 231) \\
&= \begin{pmatrix} 1 & | & 23 \\ 1 & | & 23 \end{pmatrix} + \begin{pmatrix} 2 & | & 13 \\ 1 & | & 23 \end{pmatrix} + \begin{pmatrix} 3 & | & 12 \\ 1 & | & 23 \end{pmatrix}.
\end{aligned}$$

$$\begin{aligned}
\Delta_1 \left(\begin{array}{c|c|c} 13 & 256 & 4 \\ 15 & 234 & 6 \end{array} \right) &= \Delta_1(125634 \circ Id_{\{\{1,3\}, \{2,5,6\}, \{4\}\}}) \\
&= \varepsilon \otimes 125634 \circ Id_{\{\{1,3\}, \{2,5,6\}, \{4\}\}} + 125634 \circ Id_{\{\{1,3\}, \{2,5,6\}, \{4\}\}} \otimes \varepsilon \\
&= () \otimes \begin{pmatrix} 13 & | & 256 & | & 4 \\ 15 & | & 234 & | & 6 \end{pmatrix} + \begin{pmatrix} 13 & | & 256 & | & 4 \\ 15 & | & 234 & | & 6 \end{pmatrix} \otimes ().
\end{aligned}$$

The two structures of Hopf algebra are dual to each other for the scalar product

$$\langle f, g \rangle = \delta_{f, g^{-1}}. \quad (6)$$

In other words, we have:

$$\langle \Delta f, g \otimes h \rangle = \langle f, g \bar{\sqcup} h \rangle \quad (7)$$

$$\langle \Delta_1 f, g \otimes h \rangle = \langle f, g * h \rangle. \quad (8)$$

4 Polynomial realization.

Given two alphabets $B = \{b_i; 0 \leq i\}$ and $C = \{c_i; 0 \leq i\}$, with C totally ordered, we consider an alphabet $A = \left\{ \begin{pmatrix} b_i \\ c_j \end{pmatrix}; 0 \leq i, j \right\}$ of bi-letters endowed with two relations:

- for all i and k , $\begin{pmatrix} b_i \\ c_j \end{pmatrix} \preceq \begin{pmatrix} b_k \\ c_l \end{pmatrix}$ if and only if $c_j \leq c_l$,
- for all j and l , $\begin{pmatrix} b_i \\ c_j \end{pmatrix} \equiv \begin{pmatrix} b_k \\ c_l \end{pmatrix}$ if and only if $b_i = b_k$.

The product of two biwords is the concatenation

$$\begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \begin{pmatrix} u_2 \\ v_2 \end{pmatrix} = \begin{pmatrix} u_1 u_2 \\ v_1 v_2 \end{pmatrix}.$$

We extend the definition of the standardization to biwords. Let $w = \begin{pmatrix} u \\ v \end{pmatrix}$ be a biword. The *standardized* of w is the permutation $\text{std}(v)$.

Example 11

$$\text{std} \left[\begin{pmatrix} b_1 b_1 b_3 b_1 b_2 \\ c_2 c_3 c_2 c_1 c_6 \end{pmatrix} \right] = 24315.$$

Let $f : \mathcal{A} \rightarrow \mathcal{A}'$ be a uniform block permutation. Let ξ_f be the minimum permutation (for the lexicographic order) such that $f = \xi_f \circ \text{Id}_{\mathcal{A}}$. We say that a biword w is f -compatible if

$$- \text{std}(w) = \xi_f$$

and

$$- \text{ if } i \text{ and } j \text{ are in the same block in } \mathcal{A}, \text{ then the bi-letters } w_i \text{ and } w_j \text{ satisfy } w_i \equiv w_j.$$

We set:

$$\mathbf{G}_f(A) = \sum_{w; w \text{ is } f\text{-compatible}} w. \quad (9)$$

Example 12 Let $f = \left(\begin{array}{c|cc} 13 & 256 & 4 \\ \hline 15 & 234 & 6 \end{array} \right)$ be a uniform block permutation. The permutation $\sigma = 125634$ is the minimum permutation associated with f and we have

$$\mathbf{G}_f(A) = \sum_{\substack{i,j,k \\ \text{std}(c_1 c_2 \dots c_6) = \sigma}} \begin{pmatrix} b_i b_j b_i b_k b_j b_j \\ c_1 c_2 c_3 c_4 c_5 c_6 \end{pmatrix}.$$

Theorem 1 The polynomials $\mathbf{G}_f(A)$ provide a realization of the Hopf algebra of uniform block permutations. That is to say

- $\mathbf{G}_f(A) \mathbf{G}_g(A) = \sum_{h \in f * g} \mathbf{G}_h(A)$
- Let A' be an alphabet isomorphic to the alphabet A such that, for any bi-letter w' of A' and for any bi-letter w of A , we have

$$w \preceq w'.$$

Define $A + A'$ as the disjoint union of A and A' endowed with the relations \preceq and \equiv . If we allow the bi-letters of A and A' to commute and identify $P(A)Q(A')$ with $P \otimes Q$, we have:

$$\mathbf{G}_f(A + A') = \Delta(\mathbf{G}_f). \quad (10)$$

Example 13

$$\begin{aligned} \mathbf{G} \begin{pmatrix} 1 \\ 1 \end{pmatrix} (A) \mathbf{G} \begin{pmatrix} 1 \\ 1 \end{pmatrix} (A) &= \sum_{\substack{b \in B \\ c \in C}} \binom{b}{c}^2 = \sum_{\substack{b_1, b_2 \in B \\ c_1, c_2 \in C}} \begin{pmatrix} b_1 b_2 \\ c_1 c_2 \end{pmatrix} \\ &= \sum_{\substack{b_1, b_2 \in B \\ c_1, c_2 \in C; c_1 \leq c_2}} \begin{pmatrix} b_1 b_2 \\ c_1 c_2 \end{pmatrix} + \sum_{\substack{b_1, b_2 \in B \\ c_1, c_2 \in C; c_1 > c_2}} \begin{pmatrix} b_1 b_2 \\ c_1 c_2 \end{pmatrix} \\ &= \mathbf{G} \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} (A) + \mathbf{G} \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} (A). \end{aligned}$$

$$\begin{aligned}
\mathbf{G} \left(\begin{array}{c|c} 12 & \\ \hline 12 & \end{array} \right)^{(A)} \mathbf{G} \left(\begin{array}{c} 1 \\ 1 \end{array} \right)^{(A)} &= \sum_{\substack{b \in B \\ \text{std}(v)=12}} \binom{bb}{v} \sum_{\substack{b \in B \\ c \in C}} \binom{b}{c} \\
&= \sum_{\substack{i_1, i_2 \\ \text{std}(v_1)=12 \\ \text{std}(v_2)=1}} \binom{b_{i_1} b_{i_1} b_{i_2}}{v_1 v_2} \\
&= \sum_{\substack{i_1, i_2 \\ c_1 \leq c_2 \leq c_3}} \binom{b_{i_1} b_{i_1} b_{i_2}}{c_1 c_2 c_3} + \sum_{\substack{i_1, i_2 \\ c_1 \leq c_3 < c_2}} \binom{b_{i_1} b_{i_1} b_{i_2}}{c_1 c_2 c_3} + \sum_{\substack{i_1, i_2 \\ c_3 < c_1 \leq c_2}} \binom{b_{i_1} b_{i_1} b_{i_2}}{c_1 c_2 c_3} \\
&= \mathbf{G} \left(\begin{array}{c|c} 12 & 3 \\ \hline 12 & 3 \end{array} \right) + \mathbf{G} \left(\begin{array}{c|c} 12 & 3 \\ \hline 13 & 2 \end{array} \right) + \mathbf{G} \left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right).
\end{aligned}$$

Example 14 We set $\tau = 2431$.

$$\begin{aligned}
\mathbf{G} \left(\begin{array}{c|c} 13 & 24 \\ \hline 13 & 24 \end{array} \right)^{(A + A')} &= \sum_{\substack{i, j \\ c_1 \leq c_2 \leq c_3 \leq c_4}} \binom{b_i b_j b_i b_j}{c_1 c_2 c_3 c_4} + \sum_{\substack{i, j \\ c'_1 \leq c'_2 \leq c'_3 \leq c'_4}} \binom{b'_i b'_j b'_i b'_j}{c'_1 c'_2 c'_3 c'_4} \\
&= \mathbf{G} \left(\begin{array}{c|c} 13 & 24 \\ \hline 13 & 24 \end{array} \right)^{(A)} + \mathbf{G} \left(\begin{array}{c|c} 13 & 24 \\ \hline 13 & 24 \end{array} \right)^{(A')}.
\end{aligned}$$

$$\begin{aligned}
\mathbf{G} \left(\begin{array}{c|c|c} 13 & 2 & 4 \\ \hline 23 & 4 & 1 \end{array} \right)^{(A + A')} &= \sum_{\substack{i, j, k \\ \text{std}(c_1 c_2 c_3 c_4)=\tau}} \binom{b_i b_j b_i b_k}{c_1 c_2 c_3 c_4} + \sum_{\substack{i, j, k \\ \text{std}(c'_1 c'_2 c'_3 c'_4)=\tau}} \binom{b'_i b'_j b'_i b'_k}{c'_1 c'_2 c'_3 c'_4} \\
&+ \sum_{\substack{i, j, k \\ \text{std}(c_1 c_2 c_3 c_4)=\tau}} \binom{b_i b'_j b_i b_k}{c_1 c'_2 c_3 c_4} + \sum_{\substack{i, j, k \\ \text{std}(c_1 c_2 c_3 c_4)=\tau}} \binom{b'_i b'_j b'_i b'_k}{c'_1 c'_2 c'_3 c'_4} \\
&= \mathbf{G} \left(\begin{array}{c|c|c} 13 & 2 & 4 \\ \hline 23 & 4 & 1 \end{array} \right)^{(A)} + \mathbf{G} \left(\begin{array}{c} 1 \\ 1 \end{array} \right)^{(A)} \mathbf{G} \left(\begin{array}{c|c} 13 & 2 \\ \hline 12 & 3 \end{array} \right)^{(A')} \\
&+ \mathbf{G} \left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right)^{(A)} \mathbf{G} \left(\begin{array}{c} 1 \\ 1 \end{array} \right)^{(A')} + \mathbf{G} \left(\begin{array}{c|c|c} 13 & 2 & 4 \\ \hline 23 & 4 & 1 \end{array} \right)^{(A')}.
\end{aligned}$$

5 Packed words.

We recall the definition of a packed word and the structure of Hopf algebra given in [9]. The *packed word* $\text{pack}(w)$ associated with a word $w \in A^*$ is the word u obtained by the following process. If $b_1 < b_2 < \dots < b_r$ are the letters occurring in w , u is the image of w by the homomorphism $b_i \rightarrow a_i$. A word w is said to be *packed* if $\text{pack}(w)$ is w .

5.1 WQSym*.

The Hopf algebra **WQSym*** is the \mathbb{K} -vector space generated by packed words endowed with the shifted shuffle product and the coproduct given by, using notations in [5]:

$$u \sqcup v = u \sqcup v[\max(u)], \quad (11)$$

$$\Delta(w) = \sum_{u \cdot v = w} \text{pack}(u) \otimes \text{pack}(v). \quad (12)$$

Let Ψ be the map that associates the word w of the length n with a permutation σ of size n and a set partition \mathcal{A} of $\{1, 2, \dots, n\}$ constructed by the following process. The i th letter u_i of u is the minimum of the letters σ_j whose values σ_j are in the same block in \mathcal{A} as σ_i . The word w is obtained by packing. The map Ψ induces a linear map from the vector space generated by uniform block permutations to that generated by packed words. We define it by writing uniform block permutations f in the form $Id_{\mathcal{A}'} \circ \sigma$ and applying Ψ to σ and \mathcal{A}' . We also call this map Ψ .

Example 15

$$\Psi \left[\left(\begin{array}{c|c} 12 \\ \hline 12 \end{array} \right) \right] = 11 \quad \Psi \left[\left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right) \right] = 221 \quad \Psi \left[\left(\begin{array}{c|c} 13 & 24 \\ \hline 12 & 34 \end{array} \right) \right] = 1212$$

If we endow the vector spaces with the shifted shuffle product, the map Ψ is an algebra morphism.

Proposition 1 *Let f and g be two uniform block permutations.*

$$\Psi(f \sqcup g) = \Psi(f) \sqcup \Psi(g). \quad (13)$$

Example 16

$$\begin{aligned} \Psi \left[\left(\begin{array}{c|c} 12 & 3 \\ \hline 13 & 2 \end{array} \right) \sqcup \left(\begin{array}{c|c} 1 & 2 \\ \hline 1 & 2 \end{array} \right) \right] &= 11234 + 11324 + 11342 + 13124 + 13142 + \\ &\quad 13412 + 31124 + 31142 + 31412 + 34112 \\ &= 112 \sqcup 12 \\ &= \Psi \left[\left(\begin{array}{c|c} 12 & 3 \\ \hline 13 & 2 \end{array} \right) \right] \sqcup \Psi \left[\left(\begin{array}{c|c} 1 & 2 \\ \hline 1 & 2 \end{array} \right) \right]. \end{aligned}$$

The map Ψ cannot be a coalgebra morphism as shown in the example

$$\begin{aligned} \Psi \otimes \Psi \circ \Delta \left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right) &= \Psi \otimes \Psi \left[\left(\begin{array}{c} \\ \end{array} \right) \otimes \left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right) + \left(\begin{array}{c|c} 12 & \\ \hline 12 & \end{array} \right) \otimes \left(\begin{array}{c} 1 \\ 1 \end{array} \right) \right] + \\ &\quad \Psi \otimes \Psi \left[\left(\begin{array}{c|c} 12 & 3 \\ \hline 23 & 1 \end{array} \right) \otimes \left(\begin{array}{c} \\ \end{array} \right) \right] \\ &= \varepsilon \otimes 221 + 11 \otimes 1 + 221 \otimes \varepsilon \\ &\neq \varepsilon \otimes 221 + 1 \otimes 21 + 11 \otimes 1 + 221 \otimes \varepsilon \\ &\neq \Delta(221). \end{aligned}$$

The idea is that the coproduct on packed words deconcatenates the word in all suffixes and prefixes, whereas the coproduct on uniform block permutations preserves the blocks. This suggests to define another coproduct:


$$\Delta'(w) = \sum_{\substack{u \cdot v = w \\ \text{alph}(u) \cap \text{alph}(v) = \emptyset}} \text{pack}(u) \otimes \text{pack}(v).$$

Endowed with the shifted shuffle product $\bar{\sqcup}$ and with the coproduct Δ' , the vector space generated by packed words is a combinatorial Hopf algebra denoted by \mathbf{WQSym}^{**} .

Proposition 2 *The Hopf algebras \mathbf{WQSym}^* and \mathbf{WQSym}^{**} are isomorphic.*

The morphism of Hopf algebras is given by the following process. We define the following relation on packed words of the same length. Let u and v be two packed words, the cover relation is $u \preceq v$ if and only if

- $\text{std}(u) = \text{std}(v)$
- there exists a i such that $u_k = \begin{cases} v_k & \text{if } v_k \leq i \\ v_k - 1 & \text{otherwise.} \end{cases}$



The map Γ defined as

$$\begin{aligned} \mathbf{WQSym}^* &\longrightarrow \mathbf{WQSym}^{**} \\ w &\longmapsto \sum_{u \succeq w} u \end{aligned}$$

is an isomorphism of Hopf algebras. For example, the image of a permutation is a permutation, the image of the word of length n consisting only of 1 is the sum of all the non-decreasing packed words of length n .

Example 17

$$\begin{aligned} \Gamma(11 \bar{\sqcup} 1) &= \Gamma(112 + 121 + 211) \\ &= 112 + 123 + 121 + 132 + 311 + 312 \\ &= 11 \bar{\sqcup} 1 + 12 \bar{\sqcup} 1 \\ &= \Gamma(11) \bar{\sqcup} \Gamma(1) \end{aligned}$$

$$\begin{aligned} \Gamma \otimes \Gamma(\Delta(121)) &= 121 \otimes \varepsilon + 132 \otimes \varepsilon + 12 \otimes 1 + 1 \otimes 21 + \varepsilon \otimes 121 + \varepsilon \otimes 132 \\ &= \Delta'(121) + \Delta'(132) \\ &= \Delta'(\Gamma(121)). \end{aligned}$$

Proposition 3 *The map Ψ is a surjective Hopf algebra morphism from \mathbf{UBP}^* to \mathbf{WQSym}^{**} .*

5.2 WQSym.

We denote by **WQSym** the \mathbb{K} -vector space generated by the packed words endowed with the convolution product and the coproduct given by

$$u *_W v = \sum_{w=u' \cdot v' \in PW; \text{pack}(u')=u, \text{pack}(v')=v} w, \quad (14)$$

$$\Delta(u) = \sum_{0 \leq k \leq \max(u)} u|_{\{1, \dots, k\}} \otimes \text{pack}(u|_{\{k+1, \dots, \max(u)\}}) \quad (15)$$

The dual **WQSym'** of **WQSym** is the Hopf algebra endowed with the product

$$u *_W' v = \sum_{\substack{w=u' \cdot v' \in PW \\ \text{pack}(u')=u \\ \text{pack}(v')=v \\ \text{alph}(u) \cap \text{alph}(v) = \emptyset}} w \quad (16)$$

for any packed word u and v , and the coproduct Δ .

Proposition 4 *The Hopf algebra **WQSym'** and **WQSym** are isomorphic.*

In fact, the map Γ induces a dual map. This is

$$\Gamma^* : \begin{array}{ccc} \mathbf{WQSym}' & \longrightarrow & \mathbf{WQSym} \\ w & \longmapsto & \sum_{u \preceq w} u. \end{array}$$

For example:

$$\Gamma^*(1212334) = 1212334 + 1212223 + 1212333 + 1212222.$$

This map is an isomorphism of Hopf algebras.

Example 18

$$\begin{aligned} \Gamma^*(212 *_W' 11) &= \Gamma^*(21233 + 31322 + 32311) \\ &= (21233 + 21222) + (31322 + 21211) + 32311 \\ &= 212 *_W 11 \\ &= \Gamma^*(212) *_W \Gamma^*(11) \end{aligned}$$

$$\begin{aligned} \Gamma^* \otimes \Gamma^* \circ \Delta(312341) &= \Gamma^* \otimes \Gamma^*(\varepsilon \otimes 312341 + 11 \otimes 2123 + 121 \otimes 112 + 31231 \otimes 1 + 312341 \otimes \varepsilon) \\ &= \varepsilon \otimes (312341 + 312331) + 11 \otimes (2123 + 2122) + \\ &\quad 121 \otimes (112 + 111) + 31231 \otimes 1 + (312341 + 312331) \otimes \varepsilon \\ &= \Delta(312341 + 312331) \\ &= \Delta(\Gamma^*(312341)) \end{aligned}$$

Proposition 5 *The map*

$$\begin{array}{ccc} \varphi : \mathbf{WQSym}' & \longrightarrow & \mathbf{UBP} \\ w & \longmapsto & \sum_{f; \Psi(f)=w} f \end{array}$$

is an injective morphism of Hopf algebras.

References

- [1] M. Aguiar and R.C. Orellana. The Hopf algebra of uniform block permutations. *J. Alg. Combin.*, 28(1):115–138, 2008.
- [2] M. Aguiar and F. Sottile. Structure of the Malvenuto-Reutenauer Hopf algebra of permutations. *Adv. Math.*, 191(2):225–275, 2005.
- [3] A. Connes and D. Kreimer. Hopf algebras, renormalization and noncommutative geometry. *Comm. Math. Phys.*, pages 203–242, 1998.
- [4] G. Duchamp, F. Hivert, and J.-Y. Thibon. Noncommutative symmetric functions VI: Free quasi-symmetric functions and related algebras. *Internat. J. Alg. Comput.*, 12:671–717, 2002.
- [5] L. Foissy, J.-C. Novelli, and J.-Y. Thibon. Polynomial realizations of some combinatorial Hopf algebras. [arXiv:1009.2067v1](https://arxiv.org/abs/1009.2067v1) [math.CO], 2010.
- [6] J.-L. Loday and M.-O. Ronco. Hopf algebra of the planar binary trees. *Adv. Math.*, 139:293–309, 1998.
- [7] C. Malvenuto and C. Reutenauer. Duality between quasi-symmetric functions and the Solomon descent algebra. *J. alg.*, 177:967–982, 1995.
- [8] D. Manchon. On bialgebras and Hopf algebras of oriented graphs. [arXiv:1011.3032v3](https://arxiv.org/abs/1011.3032v3) [math.CO], 2011.
- [9] J.-C. Novelli and J.-Y. Thibon. Polynomial realizations of some trialgebras. *FPSAC (San Diego)*, 2006.
- [10] J.-C. Novelli and J.-Y. Thibon. Hopf algebras and dendriform structures arising from parking functions. *Fundamenta Mathematicae*, 193:189–241, 2007.
- [11] S. Poirier and C. Reutenauer. Algèbres de Hopf de tableaux. *Ann. Sci. Math. Québec*, 19:79–90, 1995.