# Vertex-colouring edge-weightings with two edge weights

# $\begin{array}{ccc} \mbox{Mahdad Khatirinejad}^{1*} & \mbox{Reza Naserasr}^2 & \mbox{Mike Newman}^{3*} \\ & \mbox{Ben Seamone}^{2\dagger} & \mbox{Brett Stevens}^{2*} \end{array}$

<sup>1</sup> Department of Communications and Networking, Aalto University, Finland

<sup>2</sup> School of Mathematics and Statistics, Carleton University, Canada

<sup>3</sup> Department of Mathematics, University of Ottawa, Canada

received 24<sup>th</sup> July 2010, accepted 24<sup>th</sup> November 2011.

An edge-weighting vertex colouring of a graph is an edge-weight assignment such that the accumulated weights at the vertices yields a proper vertex colouring. If such an assignment from a set S exists, we say the graph is S-weight colourable. It is conjectured that every graph with no isolated edge is  $\{1, 2, 3\}$ -weight colourable.

We explore the problem of classifying those graphs which are  $\{1, 2\}$ -weight colourable. We establish that a number of classes of graphs are S-weight colourable for much more general sets S of size 2. In particular, we show that any graph having only cycles of length 0 mod 4 is S-weight colourable for most sets S of size 2. As a consequence, we classify the minimal graphs which are not  $\{1, 2\}$ -weight colourable with respect to subgraph containment. We also demonstrate techniques for constructing graphs which are not  $\{1, 2\}$ -weight colourable.

Keywords: edge weighting, graph colouring

## 1 Introduction

Let G be a simple graph and S be a set of real numbers. An S-edge-weighting of G is an assignment  $w : E(G) \to S$ . Given an S-edge-weighting, the weighted degree of a vertex v, denoted w(v), is the sum of weights of the edges incident with v. An S-edge-weighting gives a vertex colouring if the weighted degrees of adjacent vertices are different. If an S-edge-weighting vertex colouring w exists, we also call w an S-weight colouring and we say G is S-weight colourable. For a positive integer k, we say G has a k-weight colouring or G is k-weight colourable if it is S-weight colourable for every set S of size k. The most commonly studied sets S are those of the form  $\{1, \ldots, k\}$ .

**Problem 1** Given a graph G with no isolated edges, find the minimum k such that G is  $\{1, ..., k\}$ -weight colourable.

<sup>\*</sup> Partially supported by NSERC Canada.

<sup>&</sup>lt;sup>†</sup> Partially supported by Fonds québécois de la recherche sur la nature et les technologies.

 $<sup>^{\</sup>ddagger}$  Emails: mahdad.khatirinejad@tkk.fi,{bseamone,brett}@math.carleton.ca,mnewman@uottawa.ca

<sup>1365-8050 © 2012</sup> Discrete Mathematics and Theoretical Computer Science (DMTCS), Nancy, France

It is not hard to verify that  $K_4$  with a single leaf attached is  $\{1, 2\}$ -weight colourable but is not  $\{0, 1\}$ -weight colourable. It follows that the S-weight colourability of a graph is not only dependent on the size of S but also on the particular elements of S. However, if a graph G is S-weight colourable then there exists an  $i_0 = i_0(G, S)$  such that for all  $i > i_0$  the graph is also  $\{s + i : s \in S\}$ -weight colourable. One such value for  $i_0$ , though not necessarily the smallest, is  $i_0 = |S| \cdot \Delta(G) \cdot \max\{|s| : s \in S\}$ , where  $\Delta(G)$  is the maximum degree of G.

Let us start by considering the 2-weight colourability of a simple class of graphs – paths. If a and b are non-zero real numbers, then every path of length at least 2 has an  $\{a, b\}$ -weight colouring. Assigning the edge weights  $a, a, b, b, a, a, b, b, \ldots$ , beginning with one leaf of the path, gives such a colouring. However, a path has a  $\{0, a\}$ -weight colouring if and only if it is not of length 1 mod 4. The reader can easily check that paths of length 2, 3 and 4 have a  $\{0, a\}$ -weight colouring. However, if we let  $P = e_1, e_2, e_3, e_4, e_5$  be a path of length 5 (we omit vertex labels) then if  $w(e_2) = 0$  (or  $w(e_4) = 0$ ) then the ends of  $e_1$  ( $e_5$ ) will have equal weight. Thus the only way to achieve a  $\{0, a\}$ -weight colouring of P is if  $w(e_2) = w(e_4) = a$ . However, this implies that the ends of  $e_3$  will have the same weight, and hence a  $\{0, a\}$ -weight colouring cannot exist. These examples easily extend to longer paths; the details are left to the reader.

In general, it is unknown how difficult it is to decide if a given graph admits a  $\{1, 2\}$ -weight colouring, or more generally an  $\{a, b\}$ -weight colouring. As such, we present the following question:

#### **Problem 2** Is it NP-complete to decide whether a given graph is 2-weight colourable?

Returning to Problem 1, we state the following conjecture, due to Karoński, Łuczak, and Thomason [KŁT04], which motivates most of the known results on the  $\{1, \ldots, k\}$ -weight colourability of graphs.

#### **Conjecture 1.1** *Every graph with no isolated edge is* $\{1, 2, 3\}$ *-weight colourable.*

Karoński et al. [KŁT04] showed that the Conjecture 1.1 is true for 3-colourable graphs. They also proved that if S is any set of at least 183 real numbers which are linearly independent over the rational numbers then every graph with no isolated edge is S-weight colourable. Recently, Kalkowski et al. [KKP09] showed that every graph with no isolated edge is  $\{1, \ldots, k\}$ -weight colourable for k = 5. This result is an improvement on the previous bounds on k established by Addario-Berry et al. [ABDM<sup>+</sup>07], Addario-Berry et al. [ABDR08], and Wang et al. [WY08], who obtained the bounds k = 30, k = 16, and k = 13, respectively.

Our work in this paper is similarly motivated by Conjecture 1.1. However, where most others have attempted to lower the best known value of k as described above, our focus is on establishing which graphs are  $\{1, 2\}$ -weight colourable. Addario-Berry, Dalal and Reed [ABDR08] showed that asymptotically almost every graph is  $\{1, 2\}$ -weight colourable, however it is not known which ones are not. Chang et al and Lu et al ([CLWY10], [LYZ10]) have made some progress in determining which classes of graphs are  $\{1, 2\}$ -weight colourable, notably having shown that 3-connected bipartite graphs are one such class. A complete classification of such graphs would determine those graphs for which k = 3 is the smallest possible solution in Problem 1, and would reduce Conjecture 1.1 to just those graphs.

The results that follow are, for the most part, concerned with a more general problem than that of finding  $\{1, 2\}$ -weight colourings, namely that of finding  $\{a, b\}$ -weight colourings for more general values of a and b. In such cases, the existence of a  $\{1, 2\}$ -weight colouring follows as an unstated corollary. In Section 2, we establish a wide range of basic graphs which admit  $\{a, b\}$ -weight colourings. We also establish classes of graphs which do not admit  $\{a, b\}$ -weight colourings, but which do admit an  $\{a, b\}$ -edge weighting which is almost a proper colouring. These results provide building blocks for our results on the weight

colourability of bipartite graphs in Section 3 and of other general classes of graphs, particularly direct products of graphs, in Section 4. Of note, we show in Section 3 that if every cycle of G is of length  $0 \mod 4$ , then G is  $\{1, 2\}$ -weight colourable.

## 2 Building blocks: Weight colourings of basic graphs

We will use standard graph theory terminology; the reader may refer to [BM08] for clarification of any terms which are not specifically defined here.

The *length* of a path (walk) is defined to be the number of edges of the path (walk). A *thread* in a graph G is a walk connecting two vertices x and y, not necessarily distinct, such that the internal vertices are distinct from all others on the walk, all internal vertices have degree 2 in G, and  $\deg(x)$ ,  $\deg(y) \ge 3$ . If x and y are distinct, then the walk is in fact a path and in this case we may refer to the thread as an *ear*. If the condition that  $\deg(x)$ ,  $\deg(y) \ge 3$  is changed to  $\deg(x)$ ,  $\deg(y) \ge 2$  in either case, we have a *subthread* or *subear* respectively.

A *cut vertex* of a graph is one whose removal disconnects the graph. A graph is 2-connected if it has no cut vertex. A graph (not necessarily simple) is called *separable* if it can be decomposed into two nonempty subgraphs with exactly one vertex in common. A simple graph is separable if and only if it is not 2-connected. A maximal nonseparable subgraph of G is a *block* of G. Note that a block is isomorphic either to  $K_2$  or to a 2-connected graph. An *end block* of G is a block which contains at most one cut vertex of G.

A graph is *c*-colourable if the vertices can be coloured with *c* colours so that adjacent vertices get different colours.

 $K_n$  and  $C_n$ , respectively, denote the complete graph and the cycle on n vertices. The *Cartesian product* of two graphs G and H, denoted by  $G \Box H$ , is defined as the graph having vertex set  $V(G) \times V(H)$  where two vertices (u, u') and (v, v') are adjacent if and only if either u = v and u' is adjacent to v' in H or u' = v' and u is adjacent to v in G.

We present a few simple observations.

**Proposition 2.1** Let a, b, t be nonzero real numbers and G a graph. Then

- (i) G is  $\{a, b\}$ -weight colourable if and only if G is  $\{at, bt\}$ -weight colourable, and
- (ii) if G is  $\{a, b\}$ -weight colourable then G is  $\{p, q\}$ -weight colourable for any nonzero  $p, q \in \mathbb{R}$  which are linearly independent over  $\mathbb{Q}$ .

**Proof:** (i) This follows from the fact that  $w(u) \neq w(v)$  if and only if  $t \cdot w(u) \neq t \cdot w(v)$ . (ii) Note that if two adjacent vertices receive distinct linear combinations of a and b as weights, then the coefficients of these linear combinations will suffice for any two linearly independent nonzero reals.

From Proposition 2.1 we deduce the following, adopting the convention that 0 and 1 are relatively prime integers:

**Corollary 2.2** A graph G is 2-weight colourable if and only if G is  $\{a, b\}$ -weight colourable for every pair of relatively prime integers a and b.

Proposition 2.1 allows us to reduce our proofs of positive results on the existence of  $\{a, b\}$ -weight colourings of a graph to relatively prime integers. Results in which we show that G does not admit an  $\{a, b\}$ -weight colouring will not rely on such assumptions – we will prove them for all real a, b.

**Proposition 2.3** If G is d-regular and  $\{a, b\}$ -weight colourable for a fixed choice of a and b then (i) it is d-colourable, and (ii) it is 2-weight colourable.

**Proof:** (i) The weighted degree of each vertex must be a number of the form ta + (d - t)b for some  $0 \le t \le d$ , and a vertex of weighted degree da cannot be adjacent to a vertex of weighted degree db. Thus putting the vertices of weighted degree da or db in the same colour class gives a d-colouring.

(ii) In an  $\{a, b\}$ -edge weighting of a *d*-regular graph, the accumulated weight at any vertex is in a oneto-one correspondence with the number of incident edges of weight *a*. Thus if one choice of *a* and *b* gives a vertex colouring, then any other choice of *a* and *b* will as well.

**Corollary 2.4** If  $\chi(G) = \Delta(G) + 1$  or, equivalently (by Brooks theorem), if G is an odd cycle or a complete graph then G is not S-weight colourable for any set S of size 2.

Even though the complete graph is not S-weight colourable for any set of size 2, it has an S-edgeweighting that is very close to being an S-weight colouring. This specific weighting will be useful in constructing families of 2-weight colourable graphs and non-2-weight colourable graphs in Section 4.

**Lemma 2.5** Given  $n \ge 2$  and  $a \ne b \in \mathbb{R}$ , there is an  $\{a, b\}$ -edge-weighting of  $K_n$  such that the weighted degrees of all the vertices are distinct except for 2 of them. Furthermore, in any such  $\{a, b\}$ -edge-weighting, the degree sequence of the subgraph induced by the edges of weight a (as well as the subgraph induced by the edges of weight b) is either

$$(1,2,\ldots,\left\lfloor\frac{n}{2}\right\rfloor-1,\left\lfloor\frac{n}{2}\right\rfloor,\left\lfloor\frac{n}{2}\right\rfloor,\left\lfloor\frac{n}{2}\right\rfloor+1,\ldots,n-2,n-1),$$

or

$$(0,1,\ldots,\left\lceil\frac{n}{2}\right\rceil-2,\left\lceil\frac{n}{2}\right\rceil-1,\left\lceil\frac{n}{2}\right\rceil-1,\left\lceil\frac{n}{2}\right\rceil,\ldots,n-3,n-2).$$

**Proof:** We prove the first part with an explicit construction. Choose any two vertices and assign weight a to the edge joining them. Choose a new vertex and assign weight b to all the edges joining this vertex to the previous two vertices. Choose another vertex and assign weight a to all the edges joining this vertex to the previous three vertices. By repeating this process until all vertices are exhausted, we achieve the desired edge-weighting since the two vertices chosen first will have the same weight while the remainder of the graph is properly coloured. Note that we achieve the same result by swapping a and b in this argument.

We prove the second part of the lemma by induction on n. Suppose w is such an edge-weighting of  $K_n$ and let w(u) = w(v). It is easy to verify the claim for n = 2 and n = 3. If  $w(x) \notin \{(n-1)a, (n-1)b\}$ for every vertex x then w(x) can only take n - 2 values, a contradiction to the choice of w. If  $w(u) = w(v) \in \{(n-1)a, (n-1)b\}$  then by removing u and v, w induces an  $\{a, b\}$ -weight colouring of  $K_{n-2}$ , a contradiction to Corollary 2.4. Thus there exists a vertex  $x \neq u, v$  such that  $w(x) \in \{(n-1)a, (n-1)b\}$ . The claim follows by induction on  $K_n - x$ .

The following technical lemmata will be useful for the rest of the paper, since they establish useful tools for finding edge-weighting vertex colourings of graphs with specific structural properties.

**Lemma 2.6** Suppose G has a vertex v with a set of leaf neighbours L where  $|L| \ge \lceil \deg(v)/2 \rceil$ . Let  $a \ne b$  be real numbers with ab > 0. If  $G \setminus L$  is  $\{a, b\}$ -weight colourable, then so is G.

**Proof:** As mentioned, Proposition 2.1 allows us to only consider  $a, b \in \mathbb{Z}^+$ . Suppose w is an  $\{a, b\}$ -weight colouring of  $G \setminus L$ . The possible extensions of w to G give exactly |L| + 1 possible weights for v. Since v has at most |L| neighbours in  $G \setminus L$ , in at least one of the extensions, the weighted degree of v is different from the weighted degrees of the neighbours of v in  $G \setminus L$ . The weighted degree of v is also different from the weighted degrees of the neighbours of v in L, since ab > 0.

**Corollary 2.7** Every tree with at least 3 vertices is  $\{a, b\}$ -weight colourable, where  $a \neq b$  are real numbers with ab > 0.

**Proof:** The statement holds for any star,  $K_{1,n-1}$ , since the assignment of a to all edges achieves the desired result. As such the result holds for n = 3 since the unique tree on 3 vertices is a star. Let T be a tree on n vertices which is not a star and assume the result holds for any tree with fewer than n vertices. Every tree has a vertex v that has at least  $\lceil \deg(v)/2 \rceil$  leaf neighbours. Since T is not a star, removing the leaf neighbours of v gives a subtree T' on at least 3 vertices. By the induction hypothesis T' has an  $\{a, b\}$ -weight colouring. By Lemma 2.6, T does as well.

The following lemma establishes that we may contract long threads in a way that maintains weight colourability.

**Lemma 2.8** Let G be a graph,  $P = v_0, e_1, v_1, e_2, v_2, e_3, v_3, e_4, v_4, e_5, v_5$  be a subthread of G, and  $a \neq b$  be any two real numbers. Let  $G' = G/\{e_1, e_2, e_3, e_4\}$  Then,

- (i) If w is an  $\{a, b\}$ -weight colouring of G, then  $w(e_1) = w(e_5) \neq w(e_3)$ .
- (ii) If G' is  $\{a, b\}$ -weight colourable, then so is G.
- (iii) If  $\deg(v_0) = 2$  or  $\deg(v_5) = 2$ , then G is  $\{a, b\}$ -weight colourable if and only if G' is  $\{a, b\}$ -weight colourable.

**Proof:** (i) If  $w(e_1) \neq w(e_5)$  then either one of the two choices for  $w(e_3)$  results in an improper colouring at  $e_2$  or  $e_4$ . Hence  $w(e_1) = w(e_5)$  and  $w(e_3)$  must be distinct.

(ii) For convenience, we still denote the vertex obtained from the contraction by  $v_0$ . Suppose w' is an  $\{a, b\}$ -weight colouring of G'. Then  $w'(v_0) \neq w'(v_5)$ . Without loss of generality assume  $w'(v_0v_5) = a$ . Let w(e) = w'(e) for each  $e \notin \{e_1, e_2, e_3, e_4, e_5\}$ ,  $w(e_1) = w(e_5) = a$  and  $w(e_3) = b$ . There are two possibilities for the weights of  $e_2$  and  $e_4$ . Assigning  $w(e_2) = a$  and  $w(e_4) = b$  does not yield a proper vertex colouring of G if and only if either  $w(v_0) = 2a$  or  $w(v_5) = a + b$ . Similarly, defining  $w(e_2) = b$  and  $w(e_4) = a$  does not yield a proper vertex colouring of G if and only if either weighting works. If the first possibility gives  $w(v_0) = 2a$ , then the second must give  $w(v_5) = 2a$ . If the first possibility gives  $w(v_5) = a + b$ , then the second gives  $w(v_0) = a + b$ . In either case  $w(v_0) = w(v_5)$ , a contradiction.

(iii) Assume  $\deg(v_0) = 2$  and let  $e_0$  be the other edge incident with  $v_0$ . Suppose w is an  $\{a, b\}$ -weight colouring of G. By (i) we have  $w(e_0) = w(e_4)$  and  $w(e_1) = w(e_5)$ . Hence  $w(v_0) = w(v_4) \neq w(v_5)$ . Thus, by assigning the common weight of  $e_1$  and  $e_5$  to the edge  $v_0v_5$ , we get an  $\{a, b\}$ -weight colouring of G'.

The degree condition on the ends of P in Lemma 2.8 (iii) cannot be dropped. For example, by taking G to be the path of length 5, a = 1, and b = 2, Lemma 2.8 (iii) fails.

From this lemma we may deduce necessary and sufficient conditions for the existence of  $\{a, b\}$ -weight colourings of cycles.

**Proposition 2.9** Let a and b be any distinct real numbers. Then  $C_n$  is  $\{a, b\}$ -weight colourable if and only if  $n \equiv 0 \pmod{4}$ .

In lieu of a proof, we simply note that, by Lemma 2.8 (iii), the proof of this proposition may be reduced to the cases  $C_3$ ,  $C_4$ ,  $C_5$  and  $C_6$ . The details are left to the reader. There are  $\{a, b\}$ -edge weightings of other cycles of length 4k + 1, 4k + 2 and 4k + 3 which give vertex colourings with as few conflicts as possible. These results are largely technical, though not difficult to prove.

**Proposition 2.10** Let a and b be any distinct real numbers. Then  $C_{2k+1}$  has an  $\{a, b\}$ -edge weighting w such that only one edge e = uv has the property that w(u) = w(v).

**Proposition 2.11** Let a and b be any distinct real numbers. Then  $C_{4k+2}$  has an  $\{a, b\}$ -edge weighting w such that precisely two edges e = uv and e' = u'v' have the property that w(u) = w(v) and w(u') = w(v'). Furthermore,

- *the distance between e and e' is even,*
- e and e' may be chosen to be any two edges at an even distance, and
- *if*  $f_1$  *and*  $f_2$  *are the edges incident to* e*, then their weights are equal and can be chosen to be either* a *or* b *(similar for* e'*).*

We present a specific consequence of Proposition 2.11 which we will find useful.

**Proposition 2.12** Let k be an integer,  $k \ge 1$ . Then  $C_{4k+2}$  has an  $\{a, b\}$ -edge-weighting such that three consecutive vertices have equal weight and the rest of the cycle is properly coloured. Furthermore, the edge-weighting can be chosen so that the weights of the four edges which contribute to the weights of those three vertices will all be a, all b, or alternate between a and b.

Let  $\Theta_{(m_1,...,m_d)}$ ,  $d \ge 3$ , be the graph constructed from d internally disjoint paths between distinct vertices x and y, where the *i*-th path has of length  $m_i$ . For simplicity, we assume  $m_1 \le m_2 \le \cdots \le m_d$ . Such graphs will be referred to as *theta graphs*. We present necessary and sufficient conditions for theta graphs to be 2-weight colourable.

**Theorem 2.13** Let  $d \ge 3$  and let a, b be real numbers. The graph  $\Theta_{(m_1,m_2,...,m_d)}$  is 2-weight colourable if and only if it is not of the form  $\Theta_{(1,4k_2+1,...,4k_d+1)}$ .

**Proof:** Let x and y be the two vertices of degree greater than two, and let  $\{P_i | 1 \le i \le d\}$  be the d internally disjoint paths between x and y.

Suppose w is an  $\{a, b\}$ -weight colouring of  $\Theta_{(1,4k_2+1,\ldots,4k_d+1)}$ . By applying Lemma 2.8 (i) to each  $P_i$ , we observe that on any of the d disjoint paths between x and y the first and last edges must receive same weight. Thus w(x) = w(y), a contradiction since x and y are adjacent. Hence  $\Theta_{(1,4k_2+1,\ldots,4k_d+1)}$  is not  $\{a, b\}$ -weight colourable for any a, b.

Consider  $\Theta_{(m_1,m_2,...,m_d)} \ncong \Theta_{(1,4k_2+1,...,4k_d+1)}$ . We can assume that  $|a| \ge |b|$ . Let  $n_j$  be the number of paths that have length equivalent to  $j \mod 4$ . Note that  $n_0 + n_1 + n_2 + n_3 = d = \deg(x) = \deg(y)$ . For each path  $P_i$ , weight each edge according to Lemma 2.8 so that the edges incident with x are weighted

a. Then w(x) = da so it has no conflicts with its neighbours since  $d \ge 3$  and the condition on the magnitudes of a and b gives  $da \notin \{2a, a + b\}$ . Note that, if  $|P_i| \ge 2$ , there are two choices for the next edge's weight on  $P_i$  which determines the rest of the weights. Given one such weighting of a path  $P_i$ , the effects of switching to the alternate weighting where the edge incident to x receives weight a depend on the parity of the length of the path. If  $|P_i|$  is even, the weight of the edge incident to y and the vertex weights of the neighbours of x and y on  $P_i$  all change. If  $|P_i|$  odd, the weight of the edge incident to y remains unchanged, but the vertex weights of the neighbours of x and y on  $P_i$  do change. In all cases the only possible weights on path-neighbours of x or y are 2b, a + b and 2b. We prove, by cases, that there is an appropriate set of choices which make w(y) distinct from its neighbours.

 $n_0 + n_2 \ge 4$ : Our choice of weightings for even  $P_i$ 's give at least 5 possible values for w(y), so there is a choice such that  $w(y) \notin \{2b, a + b, 2a, da\}$ .

 $\mathbf{n_0} + \mathbf{n_2} = \mathbf{3}$ : If no  $P_i$  has length 1, da is not a forbidden weight for y. Also, if  $n_3 \ge 1$  then there is an edge incident to y with weight b, and  $w(y) \ne da$ . In either case there is a choice of weightings so that  $w(y) \ne \{2b, a+b, 2a\}$ .

So, assume that  $m_1 = 1$  and  $n_3 = 0$ . If the initial weighting fails then we must have

$$\{2b, a+b, 2a, da\} = \{(d-3)a+3b, (d-2)a+2b, (d-1)a+b, da\}$$

which implies that b = -(d-3)a and  $d \ge 4$ . The fact that  $|a| \ge |b|$  gives that d = 4, implying  $n_1 = 1$ and b = -a. We weight all edges explicitly. The single edge on the path of length 1 receives weight a. If  $n_0 = 3$  then weight the edges of one even path  $a, a, \ldots, -a, -a$  and the other two  $a, -a, \ldots, -a, a$ . If  $n_0 = 2$  and  $n_2 = 1$  weight the edges of the paths of length 0 mod 4 with  $a, a, \ldots, -a, -a$  and the other even path with  $a, a, \ldots, a, a$ . If  $n_0 = 1$  and  $n_2 = 2$  weight the edges of the path of length 0 mod 4 with  $a, -a, \ldots, -a, a$  and the two other even paths with  $a, -a, \ldots, a, -a$ . Finally if  $n_0 = 0$  and  $n_2 = 3$ weight the edges of all even paths with  $a, -a, \ldots, a, -a$ . Each weighting gives a vertex-colouring for its respective case.

 $n_0 + n_2 = 2$ : If  $n_3 = 0$  and  $n_0 > 0$  then assign weights to the edges of one path which is length 0 mod 4 so that the weights of the first and last edges are both a. Weight the edges of the other even path so that the edge incident to x is weighted a and the edge incident to y is weighted b. If  $n_3 = n_0 = 0$  but either d > 3 or  $b \neq 0$  then assign weights to the edges of both even paths so that their edges incident with x are weighted a, one of the edges incident with y is weighted a and the edge incident with x is weighted b. In both cases weight the edges of the paths of length 1 mod 4 so the weights are, in order beginning with the edge incident with x, a, a,  $\dots$  b, a (if the path is a single edge, give it weight a). In the case when  $n_3 = n_0 = 0$ , d = 3 and b = 0 weight the edges of the two even paths a,  $0, 0, \dots a, a, 0$  and the single odd path with  $0, 0, a, a, \dots a, 0$  (beginning with the edge incident with x in each case). The weighting given in each case gives a proper vertex colouring.

Assume  $n_3 \ge 1$ . If  $n_0 \ne n_2$  then choose weightings for each  $P_i$  so that w(x) = da and each remaining neighbour of y has accumulated weight a + b. Then  $w(y) = an_0 + an_1 + bn_2 + bn_3$ . Since  $n_3 \ge 1$  we have  $w(x) \ne w(y)$ , so the only possible conflict is if w(y) = a + b. In this case change both even  $P_i$ 's to

their alternate weighting, maintaining w(x) = da and producing a new weight at y:

$$w'(y) = bn_0 + an_1 + an_2 + bn_3$$
  
=  $w(y) + (a - b)(n_2 - n_0)$   
=  $a + b + (a - b)(n_2 - n_0)$   
=  $\begin{cases} 3a - b & \text{if } n_0 = 0, n_2 = 2\\ 3b - a & \text{if } n_0 = 2, n_2 = 0 \end{cases}$ 

In either case  $w'(y) \neq a+b$ . If  $n_0 = 2$  then y has neighbours with weights 2b, and  $3b - a \neq 2b$ . Similarly if  $n_2 = 2$  then the weight at y avoids conflict with its neighbours with weight 2a.

If  $n_0 = n_2 = 1$  we start again with choices from the basic strategy that leave all path-neighbours with weight a + b. We have  $w(x) = da \neq (n_1 + 1)a + (n_3 + 1)b = w(y)$ . Thus the only conflict can again be if w(y) = a + b or equivalently,  $an_1 + bn_3 = 0$ . In this case we weight the edges of  $P_i$ 's of lengths equivalent to 0, 1, 2, and 3 mod 4 with  $\{a, a, \ldots, b, b\}$ ,  $\{a, b, \ldots, a, a\}$ ,  $\{a, a, \ldots, a, a\}$  and  $\{a, b, \ldots, b, b\}$  respectively. We still have that  $w(y) = a + b \neq da = w(x)$  and no neighbour of y has weight a + b.

 $\mathbf{n_0} + \mathbf{n_2} = \mathbf{1}$ : If  $n_3 = 0$  then weight the edges of the even path so that the edge incident with x receives weight a and the edge incident with y receives weight b. Weight the edges of the paths of length 1 mod 4 so the weights are, in order beginning with the edge incident with x,  $a, a \dots b, a$  (if the path is a single edge, give it weight a). This weighting gives a proper vertex colouring. Assume  $n_3 \ge 1$ . Again, weight the edges of each  $P_i$  so that w(x) = da and each neighbor of y (distinct from x) has accumulated weight a + b. Since  $n_3 \ge 1$  we have that  $w(x) = da \neq w(y)$ . If  $w(y) \neq a + b$ , then w is an  $\{a, b\}$ -weight colouring. Suppose w(y) = a + b. Equivalently

$$(n_0 + n_1 - 1)a + (n_2 + n_3 - 1)b = 0.$$
(1)

Change the edge weights of the even length path to begin with b, a. Call this weighting w'. We now have w'(x) = (d-1)a + b and  $w'(y) \neq a + b$ . All neighbours of y still have weight a + b, so the only possible conflicts are between x and its neighbours. We reduce all potential conflicts to one of four cases, which are solved explicitly.

If w'(x) = w'(y) then since w'(x) = (d-1)a + b, y is incident with precisely one edge with weight b. Since  $n_3 \ge 1$ , the edge with weight b comes from a path of length 3 mod 4. This gives  $n_0 = 0$ ,  $n_2 = 1$  and  $n_3 = 1$  and then Equation 1 and  $|a| \ge |b|$  gives either

- $n_0 = 0, n_1 = 1, n_2 = 1, n_3 = 1$  and b = 0 (case iii. below).
- $n_0 = 0, n_1 = 2, n_2 = 1, n_3 = 1$  and b = -a (case iv. below).

The neighbours of x have accumulated weights either a + b or 2a. If w(x) = (d-1)a + b = a + b then this implies that d = 2 but the hypotheses of the theorem include  $d \ge 3$ . If w(x) = (d-1)a + b = 2athen b = -(d-3)a. The fact that  $d \ge 3$  and  $|a| \ge |b|$  now give either

- $n_0 = 1$ ,  $n_1 = 0$ ,  $n_2 = 0$ ,  $n_3 = 2$  and b = 0 which is dealt with in case i. below.
- $n_0 = 1$ ,  $n_1 = 1$ ,  $n_2 = 0$ ,  $n_3 = 2$  and b = -a which is dealt with in **case ii.** below.
- $n_0 = 0$ ,  $n_1 = 1$ ,  $n_2 = 1$ ,  $n_3 = 1$  and b = 0 which is dealt with in **case iii.** below.

Vertex-colouring edge-weightings with two edge weights

•  $n_0 = 0$ ,  $n_1 = 2$ ,  $n_2 = 1$ ,  $n_3 = 1$  and b = -a which is dealt with in case iv. below.

**case i.** In this case x and y are not adjacent. Weight the edges of the path of length equivalent to  $0 \mod 4$  with  $0, 0, \ldots, a, a$  and the two odd paths with  $a, 0, \ldots, 0, 0$ .

**case ii.** In this case x and y are not adjacent. Weight the edges of the paths of lengths equivalent to 0 mod 4, 1 mod 4 and 3 mod 4 with  $-a, -a, \ldots, a, a, a, -a, \ldots, a, a$  and  $a, -a, \ldots, -a, -a$  respectively.

**case iii.** In this case x and y may be adjacent. Weight the edges of the paths of lengths equivalent to  $1 \mod 4$ ,  $2 \mod 4$  and  $3 \mod 4$  with  $a, a, \ldots, 0, a, 0, 0, \ldots, 0, 0$  and  $0, 0, \ldots, 0, a$  respectively.

**case iv.** In this case x and y may be adjacent. Weight the edges of the paths of lengths equivalent to 1 mod 4, 2 mod 4 and 3 mod 4 with  $a, -a, \ldots, a, a, -a, -a, \ldots, -a, -a$  and  $a, -a, \ldots, -a, -a$  respectively.

Each of these edge-weightings gives a proper vertex colouring.

 $\mathbf{n_0} + \mathbf{n_2} = \mathbf{0}$ : Every weighting of the paths  $P_i$  which gives w(x) = da must give  $w(y) = an_1 + bn_3$ . If  $m_1 = 1$  then, since our graph is not  $\Theta_{(1,4k_2+1,\ldots,4k_d+1)}$ , we have  $n_3 \ge 1$  and thus  $w(x) \ne w(y)$ . Suppose  $m \ne 1$ . For each  $P_i$  we have two choices for y's neighbour. Each choice leaves w(y) constant. Thus there is a choice for each path which gives an edge-weighting vertex-colouring.  $\Box$ 

# 3 Bipartite graphs

We begin the section by noting that the property of being  $\{a, b\}$ -weight colourable is not one that is inherited by subgraphs, nor is the property of being non- $\{a, b\}$ -weight colourable. For example, the graph consisting of  $K_4$  with a leaf attached is  $\{1, 2\}$ -weight colourable, however  $K_4$  is not  $\{a, b\}$ -colourable for any choice of a and b. Similarly  $K_4$  contains the subgraph  $C_4$  which is 2-weight colourable.

We can, however, characterize the minimal graphs with respect to subgraph containment in the class of graph which are not  $\{a, b\}$ -weight colourable for many pairs  $\{a, b\}$  (in particular,  $\{1, 2\}$ ). In Theorem 3.9 we establish that any graph which is not  $\{a, b\}$ -weight colourable must contain  $C_{2k+1}$  or  $C_{4k+2}$  as a subgraph for some positive integer k.

**Definition 3.1** A graph G is round if every cycle of G has length 0 mod 4.

The class of round graphs is much richer than merely those obtained by taking a graph and subdividing each edge into a path of length 4. For example,  $\Theta_{(2,2,2)} \cong K_{2,3}$  is a round graph which is not obtained in this way.

The following lemma establishes a useful subgraph condition of round graphs which we will use in our study of the  $\{a, b\}$ -weight colourability of round graphs.

**Proposition 3.2** If G is a round graph and  $\Theta_{(i,j,k)}$  is a subgraph of G, then i, j and k are even and  $i \equiv j \equiv k \pmod{4}$ .

**Proof:** Let  $\Theta_{(i,j,k)}$  be a subgraph of G and let  $P_i$ ,  $P_j$  and  $P_k$  be the corresponding paths of length i, j and k respectively. Since G is round,  $i + j \equiv i + k \equiv j + k \equiv 0 \pmod{4}$ . The result follows.

Before proceeding with our results on bipartite graphs we present the following definition which we adopt throughout this section (and this section only). In any  $\{a, b\}$ -edge-weighting of a graph, the weighted degree of every vertex is of the form ra + sb for some nonnegative integers r, s. We will call a weighted vertex *even (odd)* if its weighted degree is ra + sb with r even (odd). Note that the parity

of a weighted vertex does not necessarily refer to the parity of its weight. However, by Proposition 2.1, if a and b are not independent over  $\mathbb{Q}$  then we will assume that they are relatively prime integers, and so we will assume that a is an odd integer in this case. If b is even, which will be the case in a number of the following results, then the parity of a weighted vertex does coincide with the parity of its weight.

Since a number of our arguments rely on this notion of parity, we often exclude those pairs of numbers whose ratio may be reduced to a ratio of odd integers. We define the sets:

$$\mathcal{E} = \left\{ \{a, b\} \middle| \frac{a}{b} = \frac{p}{q}, \ p, q \text{ odd integers} \right\}$$
$$\mathcal{N} = \left\{ \{a, b\} \middle| \frac{a}{b} = \frac{p}{q}, \ p, q \in \mathbb{Z}, \ pq \le 0 \right\}$$

We have already seen examples of bipartite graphs which are 2-weight colourable ( $C_{4k}$  for any  $k \ge 1$ , bipartite theta graphs except  $\Theta_{(1,4k_2+1,\ldots,4k_d+1)}$ ) and some which are not ( $C_{4k+2}$  for any  $k \ge 1$ ). From these examples, we note that a bipartite graph G with both parts of odd size is not necessarily  $\{a, b\}$ weight colourable. However, if G has one part of even size, we are able to prove G is  $\{a, b\}$ -weight colourable for particular values of a and b.

**Theorem 3.3** Let  $a, b \in \mathbb{R}$  be such that  $\{a, b\} \notin \mathcal{E}$ . If G is a connected bipartite graph with at least one part being of even size, then G is  $\{a, b\}$ -weight colourable.

**Proof:** Let  $V(G) = X \cup Y$  be a bipartition of the vertices of G with |X| even. By Corollary 2.2 and since  $\{a, b\} \notin \mathcal{E}$ , we may assume that a is an odd integer and b is an even integer. We assign the weight b to each edge of G. Clearly v is even for each  $v \in V(G)$ . Let  $V(X) = \{x_1, x_2 \dots x_{2k}\}$  and let  $P_i$  be an  $x_{2i-1}x_{2i}$ -path in G. By changing every edge weight along  $P_1$  we only change the parity of  $x_1$  and  $x_2$ . By repeating this process for each  $P_i$  we have that every vertex of X has odd parity and every vertex of Y has even parity.

Call the resulting edge-weighting w. Suppose that w is not an  $\{a, b\}$ -weight colouring. Then there are adjacent vertices x and y such that w(x) = w(y). Thus there exist integers r, r', s, s' such w(x) = ra + sb where r is odd, w(y) = r'a + s'b where r' is even, and ra + sb = r'a + s'b. If a and b are linearly independent over  $\mathbb{Q}$ , we must have r = r', a contradiction. Hence b = (p/q)a for some  $p, q \in \mathbb{Z}$  with gcd(p,q) = 1. Thus rq + sp = r'q + s'p. Since r is odd and r' is even, p even implies q must be even, a contradiction. Hence p is odd. Similarly, q is odd. Therefore, b/a = p/q with p, q odd, contradicting our choice of a and b. Thus w is an  $\{a, b\}$ -weight colouring of G.

**Corollary 3.4** Let  $a, b \in \mathbb{R}$  be such that  $\{a, b\} \notin \mathcal{E} \cup \mathcal{N}$ . Let  $G \neq K_2$  be a connected bipartite graph with a vertex of degree 1. Then G is  $\{a, b\}$ -weight colourable. In particular, trees are  $\{a, b\}$ -weight colourable.

**Proof:** Let  $V(G) = X \cup Y$  be a bipartition of the vertices of G. Let  $x \in X$  be a vertex of degree 1 and let  $y \in Y$  be its neighbour. If |X| or |Y| is even, then G is  $\{a, b\}$ -weight colourable by Theorem 3.3. If |X| is odd, then G - x has an  $\{a, b\}$ -weight colouring by Theorem 3.3, say w', such that vertices in  $X \setminus \{x\}$  are odd and vertices in Y are even. By assigning b to the edge xy we maintain the parity of all the vertices. Also, since  $\{a, b\} \notin \mathcal{N}$  we have  $w'(y) \neq 0$  and so x and y will receive different weights, thus giving an  $\{a, b\}$ -weight colouring of G.

**Theorem 3.5** Let  $a, b \in \mathbb{R}$  be such that  $\{a, b\} \notin \mathcal{E} \cup \mathcal{N}$ . Let G be a connected bipartite graph with a thread of even length P and let U be the internal vertices of P. If G - U is connected then G is  $\{a, b\}$ -weight colourable.

**Proof:** We may assume that a is a positive odd integer and b is a positive even integer. If  $X \cup Y$  is the bipartition of V(G) and either |X| or |Y| is even, then G is  $\{a, b\}$ -weight colourable by Theorem 3.3. Assume both parts of G are of odd size. Let x and y be the ends of P. We first assume that x and y are distinct. By Lemma 2.8, we may assume that P is a path of length either 2 or 4.

Consider the case that P is of length 2, say P = xvy. Let G' be the bipartite graph obtained from G by deleting v and adding two leaves,  $v_1$  adjacent to x and  $v_2$  adjacent to y. Now G' is connected and bipartite with an even side, where  $v_1$  and  $v_2$  both belong to the even side. Theorem 3.3 gives an  $\{a, b\}$ -weight colouring of G', say w', so that  $v_1$  and  $v_2$  are both odd vertices. Hence  $xv_1$  and  $yv_2$  must both receive a as their weight. Let w be an  $\{a, b\}$ -edge-weighting of G, where  $w(xv) = w'(xv_1) = a$ ,  $w(yv) = w'(yv_2) = a$  and w(e) = w'(e) for all other edges  $e \in E(G)$ . If w is not an  $\{a, b\}$ -weight colouring of G, then either w(x) = 2a or w(y) = 2a. Without loss of generality, suppose w(x) = ra + sb = 2a (a similar argument will hold for y). Since w(xv) = a and r even, we have  $r \ge 2$ . If r = 2, then sb = 0 which implies s = 0 or equivalently  $\deg_G(x) = 2$ , a contradiction. If  $r \ge 3$ , then sb < 0 which gives b < 0, a contradiction. Thus w is an  $\{a, b\}$ -weight colouring of G.

Suppose |P| = 4. Let  $P = xv_1v_2v_3y$  and let  $G' = G - v_2$ . Now G' is bipartite with an even side X', and  $x, y \in X'$ . Theorem 3.3 gives an  $\{a, b\}$ -weight colouring of G', say w', so that  $v_1$  and  $v_3$  are both even vertices. Hence  $xv_1$  and  $yv_3$  must both receive b as their weight. Let w be an  $\{a, b\}$ -edge-weighting of G, where  $w(v_1v_2) = w(v_2v_3) = a$  and w(e) = w'(e) for all other edges  $e \in E(G)$ . If w is not an  $\{a, b\}$ -weight colouring of G, then either w(x) = a+b or w(y) = a+b. Suppose w(x) = ra+sb = a+b. Then (r-1)a = -(s-1)b, and thus r is odd. Again, we have that a and b are positive integers. Thus either r-1 < 0 or s-1 < 0. However, since  $w(xv_1) = b$ , we have  $s \neq 0$ , and since r is odd,  $r \neq 0$ . Thus w is an  $\{a, b\}$ -weight colouring of G.

Now, suppose x and y are not distinct (call this vertex x). Then P is a cycle which is an end block of G and x is a cut vertex of G. Let  $z_1$  and  $z_2$  be the neighbours of x in P. Since G' = G - U is a connected bipartite graph with one part having even size, then by Theorem 3.3 there is an  $\{a, b\}$ -weight colouring of G', say w'. We give an edge weighting w'' of P as follows:

- if P has length 2 (mod 4), then by Proposition 2.12 we may define an {a, b}-weight colouring of P, w'', so that  $w''(z_1) = w''(x) = w''(z_2) = 2a$  and P is properly coloured elsewhere;
- if P has length 0 (mod 4), then by Proposition 2.9 we may define an {a, b}-weight colouring of P, w'', so that w''(x) is the larger of 2a and 2b and P is properly coloured.

Let w be the weighting obtained by combining w' and w''. Then  $w(x) > w(z_1), w(z_2)$  and x has the same parity under w as under w'. Hence the weight of x is distinct from its neighbours in G. Since all other vertices are properly coloured by w' or w'', w gives an  $\{a, b\}$ -weight colouring of G.

**Theorem 3.6** If G is a 2-connected round graph which is not a cycle then G contains at least 2 even ears.

**Proof:** 

We first claim that G contains no proper 2-connected subgraph which contains all even ears of G. Toward a contradiction suppose H is a 2-connected maximal proper subgraph of G that contains all even ears of G. There exist two vertices of H, say x and y, which are connected by a path P such that  $H \cap P = \{x, y\}$ . Since H is 2-connected, there are also 2 edge disjoint paths P' and P'' in H between x and y. Thus  $P \cup P' \cup P''$  is a theta graph, and by Lemma 3.2 P must be of even length. Since H already contains all even ears of G,  $H' = H \cup P$  must be a proper subgraph of G but H' is also 2-connected which contradicts the maximality of H.

Now, if G has no even ear, then any cycle of G is a 2-connected subgraph containing all the even ears and this is a contradiction as G is not a cycle. If G has only one ear, let T be the ear and let x and y be the two ends of T. Then there are 2 edge disjoint paths connecting x and y, one of which must be edge disjoint from T. This path together with T forms a cycle that contains all the even ears of G, a contradiction.  $\Box$ 

**Corollary 3.7** If G is a round graph and all threads of G are odd, then G has at least two leaves.

We are now able to prove that round graphs can be edge-weight vertex-coloured with most sets of size 2.

**Theorem 3.8** Every round graph is  $\{a, b\}$ -weight colourable for  $\{a, b\} \notin \mathcal{E} \cup \mathcal{N}$ .

**Proof:** Let G be a round graph. Let B be an end block with vertex of attachment v. If B is isomorphic to  $K_2$ , then G is a bipartite graph with a leaf and thus is  $\{a, b\}$ -weight colourable by Corollary 3.4. If B is a cycle, then B is an even thread and G is  $\{a, b\}$ -weight colourable by Theorem 3.5. Otherwise, if B is a 2-connected graph which is not a cycle, then by Theorem 3.6, B has at least two even ears and thus B has at least one even ear, say P, which does not contain v as an internal vertex. Let U be the internal vertices of P. Since G - U is connected, G is  $\{a, b\}$ -weight colourable by Theorem 3.5.

Theorem 3.8, together with Proposition 2.9, gives a class of minimal subgraphs with respect to containment which cannot be  $\{a, b\}$ -weight coloured for the pairs  $\{a, b\}$  on which we have focused.

**Corollary 3.9** Let a and b be real numbers such that  $\{a, b\} \notin \mathcal{E} \cup \mathcal{N}$ . Any graph which is not  $\{a, b\}$ -weight colourable must contain a cycle of length 1, 2 or 3 mod 4.

We end this section with the following problem.

**Problem 3** Is it true that all bipartite graphs except  $C_{4k+2}$  and  $\Theta_{(1,4k_1+1,4k_2+1,...,4k_d+1)}$  are 2-weight colourable?

#### 4 More families of graphs with determined 2-weight colourability

We have given a number of examples of  $\{a, b\}$ -weight colourable graphs for values of a and b subject to particular restrictions. However we have seen few examples of graphs for which a and b can be any distinct real numbers. We note that the Petersen graph provides such an example of a 2-weight colourable graph. One such edge-weighting is given in Figure 1. By Proposition 2.3, note that any 2-weight colouring of the Petersen graph gives a 3-colouring of it, which is also an optimal proper vertex colouring.

In the rest of this section we describe more families of 2-weight colourable graphs as well as a class of nonbipartite graphs which are  $\{a, b\}$ -weight colourable when ab > 0. In particular we show that all




Fig. 1: An  $\{a, b\}$ -weight colouring of the Petersen graph. Bold edges are to receive weight b.

unicyclic graphs except cycles of length 1, 2, 3 mod 4 are 2-weight colourable. We also provide a number of results on Cartesian products of graphs. Finally, we explore techniques for constructing graphs which do not admit  $\{a, b\}$ -weight colourings for any choice of a and b.

We begin with our result on unicyclic graphs.

**Theorem 4.1** Every connected unicyclic graph except  $C_{2k+1}$  and  $C_{4m+2}$  is  $\{a, b\}$ -weight colourable, where a and b are real numbers with ab > 0.

**Proof:** We may assume that 0 < a < b. By contradiction, let G be the smallest counterexample to our claim. Let C be the only cycle of G. We first note that by Lemma 2.6, we may assume that every vertex of G is either on C or is adjacent to a vertex of C. We may also assume that every vertex of G has degree at most 3. Next, we claim that there are at least two vertices of degree at least 3 on C. If not, let v be the only vertex of degree at least 3 on C. Let x and y be the neighbours of v on C. It is easy to find an edge-weighting w of C which yields a proper colouring on C - v and  $w(v) \ge w(x), w(y)$ . By assigning b to the other edge incident with v, we get an  $\{a, b\}$ -weight colouring of G, a contradiction.

Next, we claim that G has at most one ear of length at least 2. If not, then we choose some maximal path of degree 3 vertices on  $C, x_1, \ldots, x_k$ , and remove all leaves of G adjacent to those vertices. Call this subgraph G'. By minimality of our choice of G, we can assign an  $\{a, b\}$ -weight colouring w' to G'. Let w be the weighting of E(G) given by w(e) = w'(e) if  $e \in E(G')$  and w(e) = b otherwise. The only possible conflicts are between  $x_1$  and its neighbour on C which is not  $x_2$ , say y (or, similarly, between  $x_k$  and it's neighbour on C which is not  $x_{k-1}$ ). However, since  $w(x_1) \ge a + b + w(x_1y)$  and  $w(y) \le w(x_1y) + b$  (similar for  $x_k$ ), w is an  $\{a, b\}$ -weight colouring of G which contradicts our choice of G.

If G has exactly one ear of length at least 2, let e = rs and e' = r's' be the two edges that have exactly one endpoint of degree 2. Specifically, let  $\deg(r) = \deg(r') = 2$ ,  $\deg(s) = \deg(s') = 3$ . Note that r and r' need not be distinct, but, since there are at least 2 vertices on C of degree 3, s and s' are distinct. We construct an  $\{a, b\}$ -weight colouring of G based on the length of C mod 4.

Suppose |C| is odd. By Proposition 2.10, C has an {a,b}-edge weighting w' which gives a proper vertex colouring except across rs. Let w(e) = w'(e) if e ∈ E(C). If w'(r') - w'(s') = a, let w(e) = b for all e ∈ E(G) \ E(C). Otherwise, let w(e) = a for all e ∈ E(G) \ E(C). Clearly each leaf's neighbour has a weight strictly greater than its own. Since w' gives a proper colouring of C except for r and s, the only adjacent vertices of G which might not be properly coloured are r

and s or r' and s'. However, our choice of weights for the leaves of G guarantees that r, s, r', s' are properly coloured as well. Thus w is an  $\{a, b\}$ -weight colouring of G.

- Suppose  $|C| \equiv 0 \pmod{4}$ . By Proposition 2.9, C has an  $\{a, b\}$ -weight colouring w' such that w(r) = 2a and w(s) = a + b. Let w(e) = w'(e) if  $e \in E(C)$ . If w'(r') w'(s') = a, let w(e) = b for all  $e \in E(G) \setminus E(C)$ . Otherwise, let w(e) = a for all  $e \in E(G) \setminus E(C)$ . By the same argument as above, w is an  $\{a, b\}$ -weight colouring of G.
- Suppose |C| ≡ 2 (mod 4). Let t be the other neighbour of s on C and let t' be the other neighbour of r' on C. By Proposition 2.11, there is an {a, b}-edge weighting such that all vertices are properly coloured except r, s and t, and such that w(t'r') = w(r's') = a. Let w(e) = w'(e) for all e ∈ E(C). Let f be the edge between s and its leaf, and let w(f) = a. For each e ∈ E(G) \E(C) \{f}, let w(e) = b. The only possible improperly coloured pairs of vertices are r and s, s and t or r' and s'. However,

$$\begin{array}{lll} w(r) &=& w'(r) = w'(s) < w(s) \\ w(s) &=& w'(s) + a = w'(t) + a < w'(t) + b = w(t) \\ w(r') &=& 2a < a + 2b = w(s') \end{array}$$

and so w is an  $\{a, b\}$ -weight colouring of G.

The only remaining case is that every vertex of C has degree 3. If |C| is even, assign the same weight to all the edges on the cycle and alternating weights to the leaf edges. The reader can verify that a solution for the cases when |C| = 3 or |C| = 5 exists. Each of these cases can be extended to larger odd cycle by making the replacement indicated in Figure 2. Note that the variables  $\overline{\ell}$  and  $\overline{n}$  refer to the weights different from  $\ell$  and n, respectively.




Fig. 2: Replacement operation to expand 2-weight colourings to larger cycles.

Thus, no minimal counterexample G exists.

**Proposition 4.2** For  $n \ge 4$ , the graph  $K_2 \square K_n$  is 2-weight colourable.


**Proof:** Let  $K_n$  and  $K'_n$  be the two copies of the complete graph. Denote the vertices of  $K_n$  and  $K'_n$ , respectively by

$$\{ u_1, u_2, \dots, u_{\lfloor n/2 \rfloor}, v_{\lfloor n/2 \rfloor}, v_{\lfloor n/2 \rfloor}, u_{\lfloor n/2 \rfloor+1}, \dots, v_{n-2}, v_{n-1} \}, \\ \{ u'_1, u'_2, \dots, u'_{\lfloor n/2 \rfloor}, v'_{\lfloor n/2 \rfloor}, v'_{\lfloor n/2 \rfloor+1}, \dots, v'_{n-2}, v'_{n-1} \}.$$

Let p be a derangement (permutation with no fixed points) of  $\{1, \lfloor n/2 \rfloor\}$  and  $\pi$  be a derangement of  $\{\lfloor n/2 \rfloor, n-1\}$ . Let  $u_i$  be adjacent to  $u'_{p(i)}$  for all  $1 \leq i \leq \lfloor n/2 \rfloor$  and  $v_i$  be adjacent to  $v'_{\pi(i)}$  for  $\lfloor n/2 \rfloor \leq i \leq n-1$ .

Since the graph is *n*-regular, if adjacent vertices have distinct weights then they have distinct numbers of incident edges having weight *b*. Using Lemma 2.5, we may weight the edges of  $K_n$  and  $K'_n$  so that the subscript of the vertex is precisely equal to the number of edges weighted *b* incident to that edge in  $K_n$ . Label  $u_i u'_{p(i)}$  with *a* for all  $1 \le i \le \lfloor n/2 \rfloor$  and weight  $v_i v'_{\pi(i)}$  with *b* for  $\lfloor n/2 \rfloor \le i \le n-1$ . Then any two vertices that are adjacent have a distinct number of incident edges weighted *b* and thus  $K_2 \square K_n$  is 2-weight colourable.

Figure 3 gives an illustration of this construction.



**Fig. 3:** An  $\{a, b\}$ -weight colouring of  $K_2 \square K_n$ . Bold edges are to receive weight b.


**Proposition 4.3** The graph  $K_2 \square C_n$  is 2-weight colourable if and only if  $n \ge 4$  and  $n \ne 5$ .

**Proof:** If n is even, then give every edge of one copy of  $C_n$  weight a and every edge of the other copy weight b. By alternating the weights of the images of  $K_2$  between a and b along the cycles, we have the desired  $\{a, b\}$ -weight colouring.


An example of an  $\{a, b\}$ -weight colouring of  $K_2 \square C_7$  is given in Figure 4. It can be extended to an  $\{a, b\}$ -weight colouring of  $K_2 \square C_9$  by replacing the left subgraph in Figure 5 with the right graph. Note that the right subgraph contains the left one, and thus this operation may be repeated as many times as needed to give an  $\{a, b\}$ -weight colouring for any  $K_2 \square C_{2k+1} (k \ge 3)$  The reader may verify that no  $\{a, b\}$ -weight colouring of  $K_2 \square C_3$  or  $K_2 \square C_5$  exists.  $\square$ 

**Theorem 4.4** Let G be a graph and H be a regular bipartite graph. If  $G \square K_2$  is 2-weight colourable, then  $G \square H$  is 2-weight colourable.

**Proof:** Let w be an  $\{a, b\}$ -weight colouring of  $G \square K_2$ . Denote the two copies of G by  $G_1$  and  $G_2$  and denote the vertices of  $K_2$  by  $t_1$  and  $t_2$ . Since H is regular (say *d*-regular) and bipartite, Hall's Theorem guarantees a perfect matching M of H. Let X and Y be the parts of V(H).



**Fig. 4:** An  $\{a, b\}$ -weight colouring of  $K_2 \square C_7$ . Bold edges are to receive weight b.



**Fig. 5:** Replacement operation for obtaining an  $\{a, b\}$ -weight colouring of  $K_2 \square C_{2k+1}$  for  $k \ge 4$ .

Define an edge-weighting of  $G \square H$  as follows. For each edge  $e = xy \in M$  where  $x \in X$  and  $y \in Y$ , weight the edges of the subgraph  $G \square e$  by w so that each vertex  $(u_G, x) \in V(G \square H)$  has weight  $w(u_G, t_1)$  and  $(u_G, y) \in V(G \square H)$  has weight  $w(u_G, t_2)$ . Assign every other edge of  $G \square H$  weight a. Call this weighting  $\phi$ .

We have that  $\phi(u) = w(u_G, t_1) + (d-1)a$  if  $u_H \in X$  and  $\phi(u) = w(u_G, t_2) + (d-1)a$  if  $u_H \in Y$ . Two vertices are adjacent if either their *H*-coordinates agree and they are adjacent in a copy of *G* or if their *G*-coordinates agree and they are adjacent in a copy of *H*. In the former case, their weights are distinct under  $\phi$  since they are distinct under *w*. In the latter, consider two adjacent vertices  $u = (u_G, u_H)$  and  $u' = (u_G, u'_H)$  where  $u_H \in X, u'_H \in Y$ . Then,  $w(u_G, t_1) \neq w(u_G, t_2)$  by choice of *w*, which implies that  $\phi(u) \neq \phi(u')$ . Thus  $\phi$  is an  $\{a, b\}$ -weight colouring of  $G \square H$ .

**Corollary 4.5** *If G and H are regular bipartite graphs, then the following graphs are 2-weight colourable:* 

- (i)  $K_n \Box H$ , if  $n \ge 4$
- (ii)  $C_n \Box H$  if  $n \ge 4, n \ne 5$
- (*iii*)  $G \Box H$

Vertex-colouring edge-weightings with two edge weights

**Proof:** Applying Theorem 4.4 to Propositions 4.2 and 4.3 immediately gives results (i) and (ii) respectively. For (iii), since  $K_2 \square K_2 \cong C_4$ ,  $K_2 \square K_2$  is 2-weight colourable by Proposition 2.9. By Theorem 4.4,  $K_2 \square H$  is 2-weight colourable; applying Theorem 4.4 again gives us that  $G \square H$  is 2-weight colourable.

In order to construct non-2-weight colourable graphs below, we make use of a class of "gadget" graphs. These gadgets are themselves 2-weight colourable, but they have the property that in any of their 2-weight colourings, certain edges receive a predetermined weight.

Define the graph  $\hat{K}_n$  to be the graph obtained from  $K_n$  by subdividing one edge exactly once.

**Proposition 4.6** For  $n \ge 4$ , the graph  $\hat{K}_n$  is 2-weight colourable. Moreover, in any 2-weight colouring of  $\hat{K}_n$ , the edges incident to its degree 2 vertex must receive the same colour.

**Proof:** Let x be the vertex of  $\widehat{K}_n$  of degree 2 and let u, v be its neighbours. An  $\{a, b\}$ -weight colouring of  $\widehat{K}_4$  is given in Figure 6. So assume  $n \ge 5$ . Let  $K_n$  be obtained by adding the edge uv to  $\widehat{K}_n - x$ . By Lemma 2.5, there exists an edge-weighting w of  $K_n$  such that all the vertices have distinct weighted degrees except for u and v. Moreover, w(u) = w(v) = ra + (n-1-r)b, where  $r \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil - 1\}$ . Assign the weight w(uv) from  $K_n$  to the edges xu and xv in  $\widehat{K}_n$ . Note that w is an  $\{a, b\}$ -weight colouring as long as  $w(u) = w(v) \neq w(x)$ . We have  $w(x) \in \{2a, 2b\}$ . Since a and b may be swapped in Lemma 2.5, we assume that w(x) = 2a. If  $w(u) \neq 2a$ , we are done. Suppose w(u) = 2a. We consider two cases:

- If n is odd, then the edge weighting w' given by swapping every edge's weight gives w'(u) = w(u) = 2a ≠ 2b = w'(x).
- If n is even then, by the construction of the weighting in Lemma 2.5, w(u) = n/2 a + (n/2 1)b. So, 2a = n/2 (a+b) − b. If the edge weighting w' given by swapping every edge's weight gives a conflict between u and x, then 2b = n/2 (a + b) − a. Together, these imply that a = b, a contradiction.

Thus  $\hat{K}_n$  admits an  $\{a, b\}$ -weight colouring.

To prove the second part, toward a contradiction, suppose  $\widehat{K}_n$  is the smallest counterexample for which there exists an  $\{a, b\}$ -weight colouring w such that  $w(xu) \neq w(xv)$ . By inspection, we may check that  $\widehat{K}_4$  does not admit such edge-weighting. So assume  $n \geq 5$ . Note that there exists no vertex  $y \neq u, v$  such that  $w(y) \in \{(n-1)a, (n-1)b\}$ , otherwise  $\widehat{K}_n - y$  would be a smaller counterexample. Therefore, since w induces a vertex colouring and all the weighted degrees (except for x) are of the form ra + (n-1-r)bfor some  $0 \leq r \leq n-1$ , we must have  $w(u), w(v) \in \{(n-1)a, (n-1)b\}$ . But then by removing u, v, and x we get an  $\{a, b\}$ -weight colouring of  $K_{n-2}$ , a contradiction to Corollary 2.4.

**Corollary 4.7** Given a graph G, let G' be obtained from identifying a vertex of G with the degree 2 vertex of  $\hat{K}_n$ . Then in any 2-weight colouring of G', edges in  $\hat{K}_n$  incident to its degree 2 vertex must receive same colour.

**Proof:** Since the proof of Proposition 4.6 did not depend in any way on the accumulated weight at vertex x, then regardless of graph joined to  $\hat{K}_n$  at x, the two edges incident with x in  $\hat{K}_n$  must still receive the same weight.

An example is given on the left of Figure 6. In the case  $G = K_2$  and n = 4, the weight of the leaf's edge is forced to be equal to that of its incident edges; this is another useful gadget. It is shown on the right of Figure 6.




Fig. 6: The graphs  $\hat{K}_4$  and  $\hat{K}_4$  with a leaf are 2-weight colourable. Bold edges represent one weight-class.

We use Proposition 4.6, which established the weight colourability of  $\hat{K}_n$ , to construct the following examples of non 2-weight colourable graphs.

**Example 4.8** The following graphs cannot be 2-weight coloured:

- (i) Join two copies of  $\hat{K}_4$  by an edge attached at their vertices of degree 2.
- (ii) Join 2n + 1 copies of  $\hat{K}_4$  to a  $C_{2n+1}$  by an edge attaching the degree 2 vertex in each copy of  $\hat{K}_4$  to a distinct cycle vertex.

To see why the graph defined in (ii), which we denote H, cannot be S-weight coloured for any set S of size 2, consider the accumulated weight at one of the cycle vertices, say v. Since H is 3-regular graph,  $w(v) \in \{3a, 2a+b, a+2b, 3b\}$ . If w(v) = 3a, then the noncycle edge, e, incident with v must have weight a and, as shown in Figure 6, so must the two edges in the copy of  $\hat{K}_4$  joined to v by an edge. Thus both endpoints of e would have weight 3a. A similar argument shows that  $w(v) \neq 3b$ . Thus the only possible accumulated weights on cycle vertices are 2a + b and 2b + a. Since an odd cycle cannot be properly 2-coloured, we see that H cannot be 2-weight coloured.

Our next family of gadget graphs are described below.

**Proposition 4.9** Let  $0 \le a \in \mathbb{Z}$  and  $d \mid a$ . Let H be a graph and G be a graph obtained from identifying a vertex u of H with a vertex of a  $K_n$  (all other vertices of H and  $K_n$  being disjoint). If

$$\deg_H(u) < \left(\frac{d}{a+d}\right) \left\lfloor \frac{n-1}{2} \right\rfloor$$

then G is not  $\{a, a + d\}$ -weight colourable. Furthermore, if

$$\deg_H(u) = \left(\frac{d}{a+d}\right) \left\lfloor \frac{n-1}{2} \right\rfloor,$$

then in any  $\{a, a + d\}$ -weight colouring of G, all edges in H incident to u must receive weight a + d.

**Proof:** We first prove the statement for d = 1. Toward a contradiction, suppose w is an  $\{a, a + 1\}$ -weighting colouring of G. Every vertex of  $K_n - u$  has weighted degree ra + (n - 1 - r)(a + 1) = (n - 1)(a + 1) - r for some  $0 \le r \le n - 1$  and both of the weights (n - 1)a and (n - 1)(a + 1) cannot

appear simultaneously on  $K_n - u$ . If w(u) < (n-1)(a+1) then there are only n-1 colours available for the vertices of  $K_n$ , a contradiction. So  $w(u) \ge (n-1)(a+1)$ .

Let  $w|_{K_n}$  be the edge-weighting of  $K_n$  induced by w. By Corollary 2.4,  $K_n$  is not  $\{a, a + 1\}$ -edge-weight colourable. Thus, there must be exactly two vertices of  $K_n$  with the same weight given by  $w|_{K_n}$  and u must be one such vertex. By Lemma 2.5, we get  $w|_{K_n}(u) = ra + (n - 1 - r)(a + 1) = (n - 1)(a + 1) - r$ , where  $r \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil - 1\}$ . Note that  $r \ge \lfloor (n - 1)/2 \rfloor$ . If u is incident with s edges of weight a in H, then we have

 $(n-1)(a+1) \le w(u) = ra + (n-1-r)(a+1) + sa + (\deg_H(u) - s)(a+1),$ 

which simplifies to  $\deg_H(u) \ge (\frac{1}{a+1})(r+s)$ . Hence

$$\deg_H(u) \geq \left(\frac{1}{a+1}\right) \left( \left\lfloor \frac{n-1}{2} \right\rfloor + s \right).$$

This is a contradiction since  $\deg_H(u) < (\frac{1}{a+1}) \lfloor \frac{n-1}{2} \rfloor$ . Also, if  $\deg_H(u) = (\frac{1}{a+1}) \lfloor \frac{n-1}{2} \rfloor$  then we must have s = 0, proving the second claim.

Now, let d be any positive divisor of a. By Proposition 2.1, G has an  $\{a, a + d\}$ -weight colouring if and only if it has a  $\{\frac{a}{d}, \frac{a}{d} + 1\}$ -weight colouring. If  $\deg_H(u) < \left(\frac{1}{a/d+1}\right) \lfloor \frac{n-1}{2} \rfloor$  then G has no  $\{\frac{a}{d}, \frac{a}{d} + 1\}$ -weight colouring by the above argument. Hence, if  $\deg_H(u) < \left(\frac{d}{a+d}\right) \lfloor \frac{n-1}{2} \rfloor$  then G has no  $\{a, a + d\}$ -weight colouring. The second result follows similarly.  $\Box$ 

We may use Proposition 4.9 to construct many graphs which are not  $\{a, a + 1\}$ -weight colourable and so, in particular, are not  $\{1, 2\}$ -weight colourable. In fact, if H is any graph and u any vertex of H, then there is an n large enough so that attaching  $K_n$  to u (and only to u) gives a graph which is not  $\{a, a + 1\}$ -weight colourable. We can also use the equality condition to construct graphs with no  $\{a, a + 1\}$ -weight colouring (for a specific a). For example, the graph obtained by joining two copies of  $K_n$ ,  $n \ge 5$ , with a path of length 3, say  $e_1, e_2, e_3$ , is not  $\{a, a + 1\}$ -weight colourable for  $a = \lfloor \frac{n-1}{2} \rfloor - 1$  since the weights of  $e_1$  and  $e_3$  are forced to be a + 1, and thus the ends of  $e_2$  will receive the same weight.

We finish this section with a problem whose solution would be useful in the study of  $\{1, 2, 3\}$ -weight colourable graphs.

#### Problem 4 Does there exist a graph which is either

- uniquely  $\{1, 2, 3\}$ -weight colourable (up to isomorphism), or
- {1,2,3}-weight colourable where any such colouring forces certain edges of G to receive a particular weight?

Moreover, does there exist a graph with either of these properties which maintains that property when it is attached in some way to another graph?

#### References

- [ABDM<sup>+</sup>07] Louigi Addario-Berry, Ketan Dalal, Colin McDiarmid, Bruce A. Reed, and Andrew Thomason. Vertex-colouring edge-weightings. *Combinatorica*, 27(1):1–12, 2007.
- [ABDR08] L. Addario-Berry, K. Dalal, and B. A. Reed. Degree constrained subgraphs. *Discrete Appl. Math.*, 156(7):1168–1174, 2008.
- [BM08] Adrian Bondy and U.S.R. Murty. "Graph Theory" Springer, 2008.
- [CLWY10] Gerard Chang, Changhong Lu, Jiaojiao Wu, and Qinglin Yu Vertex-coloring edgeweightings of graphs. *Taiwanese J. Math.*, 15(4):1807–1813, 2011
- [KKP09] Maciej Kalkowski, Michal Karonski, and Florian Pfender. Vertex-coloring edgeweightings: Towards the 1-2-3-conjecture. J. Combin. Theory Ser. B, 100(3):347–349, 2010.
- [KŁT04] Michał Karoński, Tomasz Łuczak, and Andrew Thomason. Edge weights and vertex colours. J. Combin. Theory Ser. B, 91(1):151–157, 2004.
- [LYZ10] Hongliang Lu, Qinglin Yu and Cun-Quan Zhang. Vertex-Coloring 2-Edge-Weightings of Graphs. *European J. Combin.*, 32(1):21–27, 2011.
- [WY08] Tao Wang and Qinglin Yu. On vertex-coloring 13-edge-weighting. *Front. Math. China*, 3(4):581–587, 2008.