Discrete Mathematics and Theoretical Computer Science DMTCS vol. **14:1**, 2012, 43–54

The generalized 3-connectivity of Cartesian product graphs[†]

Hengzhe Li[‡] Xueliang Li Yuefang Sun

Center for Combinatorics and LPMC-TJKLC, Nankai University, Tianjin, China

received 4th May 2011, revised 20th November 2011, 11th January 2012, , accepted 19th February 2012.

The generalized connectivity of a graph, which was introduced by Chartrand et al. in 1984, is a generalization of the concept of vertex connectivity. Let S be a nonempty set of vertices of G, a collection $\{T_1, T_2, \ldots, T_r\}$ of trees in G is said to be internally disjoint trees connecting S if $E(T_i) \cap E(T_j) = \emptyset$ and $V(T_i) \cap V(T_j) = S$ for any pair of distinct integers i, j, where $1 \leq i, j \leq r$. For an integer k with $2 \leq k \leq n$, the k-connectivity $\kappa_k(G)$ of G is the greatest positive integer r for which G contains at least r internally disjoint trees connecting S for any set S of k vertices of G. Obviously, $\kappa_2(G) = \kappa(G)$ is the connectivity of G. Sabidussi's Theorem showed that $\kappa(G \,\Box\, H) \geq \kappa(G) + \kappa(H)$ for any two connected graphs G and H. In this paper, we prove that for any two connected graphs G and H with $\kappa_3(G) \geq \kappa_3(H)$, if $\kappa(G) > \kappa_3(G)$, then $\kappa_3(G \,\Box\, H) \geq \kappa_3(G) + \kappa_3(H)$; if $\kappa(G) = \kappa_3(G)$, then $\kappa_3(G \,\Box\, H) \geq \kappa_3(G) + \kappa_3(H) - 1$. Our result could be seen as an extension of Sabidussi's Theorem. Moreover, all the bounds are sharp.

Keywords: Connectivity, Generalized connectivity, Internally disjoint path, Internally disjoint trees.

1 Introduction

All graphs in this paper are undirected, finite and simple. We refer to the book [1] for graph theoretic notations and terminologies not described here. For any graph G, the connectivity $\kappa(G)$ of a graph G is defined as $min\{|S| \ : S \subseteq V(G)$ and $G - S$ is disconnected or trivial$\}$. Whitney [14] showed an equivalent definition of the connectivity of a graph. For each pair of vertices x, y of G, let $\kappa(x, y)$ denote the maximum number of internally disjoint paths connecting x and y in G. Then the connectivity $\kappa(G)$ of G is $min\{\kappa(x, y) \, : \, x, y$ are distinct vertices of $G\}$.

The Cartesian product of graphs is an important method to construct a bigger graph, and plays a key role in design and analysis of networks. In the past several decades, many authors have studied the (edge) connectivity of the Cartesian product graphs. Specially, Sabidussi in [11] derived the following prefect and well-known theorem on the connectivity of Cartesian product graphs.

[†]Supported by NSFC No. 11071130.

[‡]Email: `lhz2010@mail.nankai.edu.cn`

Theorem 1.1 (Sabidussi's Theorem [11]) *Let G and H be two connected graphs. Then*

$$\kappa(G \,\square\, H) \geq \kappa(G) + \kappa(H).$$

More information about the (edge) connectivity of the Cartesian product graphs can be found in [4, 5, 6, 11, 12, 15, 16].

The generalized connectivity of a graph G, which was introduced by Chartrand et al. in [2], is a natural and nice generalization of the concept of vertex connectivity. A tree T is called an *S-tree* (*$\{u_1, u_2, \ldots, u_k\}$-tree*) if $S \subseteq V(T)$, where $S = \{u_1, u_2, \ldots, u_k\} \in V(G)$. A family of trees $T_1, T_2, \ldots, T_r$ are *internally disjoint S-trees* if $E(T_i) \cap E(T_j) = \emptyset$ and $V(T_i) \cap V(T_j) = S$ for any pair of integers i and j, where $1 \leq i < j \leq r$. We use $\kappa(S)$ to denote the greatest number of internally disjoint S-trees. For an integer k with $2 \leq k \leq n$, the *k-connectivity* $\kappa_k(G)$ of G is defined as $min\{\kappa(S) \,|\, S \in V(G) \ and \ |S| = k\}$. Clearly, when $|S| = 2$, $\kappa_2(G)$ is nothing new but the connectivity $\kappa(G)$ of G, that is, $\kappa_2(G) = \kappa(G)$, which is the reason why one addresses $\kappa_k(G)$ as the generalized connectivity of G. By convention, for a connected graph G with less than k vertices, we set $\kappa_k(G) = 1$. For any graph G, clearly, $\kappa(G) \geq 1$ if and only if $\kappa_k(G) \geq 1$.

In addition to being a natural combinatorial measure, the generalized connectivity can be motivated by its interesting interpretation in practice. For example, suppose that G represents a network. If one considers to connect a pair of vertices of G, then a path is used to connect them. However, if one wants to connect a set S of vertices of G with $|S| \geq 3$, then a tree has to be used to connect them. This kind of tree with minimum order for connecting a set of vertices is usually called a Steiner tree, and popularly used in the physical design of VLSI, see [13]. Usually, one wants to consider how tough a network can be, for the connection of a set of vertices. Then, the number of totally independent ways to connect them is a measure for this purpose. The generalized k-connectivity can serve for measuring the capability of a network G to connect any k vertices in G.

In [8], Li and Li investigated the complexity of determining the generalized connectivity and derived that for any fixed integer $k \geq 2$, given a graph G and a subset S of $V(G)$, deciding whether there are k internally disjoint trees connecting S, namely deciding whether $\kappa(S) \geq k$, is NP-complete. The generalized connectivity of complete bipartite graphs was studied by Okamoto and Zhang in [10], and Li and Li in [7].

Chartrand et al. [3] got the following result for complete graphs.

Theorem 1.2 *[3] For every two integers n and k with $2 \leq k \leq n$, $\kappa_k(K_n) = n - \lceil k/2 \rceil$.*

Theorem 1.3 *[9] Let G be a connected graph with at least three vertices. If G has two adjacent vertices with minimum degree δ, then $\kappa_3(G) \leq \delta - 1$.*

Theorem 1.4 *[9] For any connected graph G, $\kappa_3(G) \leq \kappa(G)$. Moreover, the upper bound is sharp.*

In this paper, we study the 3-connectivity of Cartesian product graphs and get the following result.

Theorem 1.5 *Let G and H be connected graphs such that $\kappa_3(G) \geq \kappa_3(H)$. The following assertions hold:*

(i) *If $\kappa(G) = \kappa_3(G)$, then $\kappa_3(G \,\square\, H) \geq \kappa_3(G) + \kappa_3(H) - 1$. Moreover, the bound is sharp;*

(ii) *If $\kappa(G) > \kappa_3(G)$, then $\kappa_3(G \,\square\, H) \geq \kappa_3(G) + \kappa_3(H)$. Moreover, the bound is sharp.*

The paper is organized as follows. In Section 2, we recall the definition and properties of Cartesian product graphs, and give some basic results about the internally disjoint S-trees. As usual, in order to get a general result, we begin with a special case. In Section 3, we study the 3-connectivity of the Cartesian product of a graph G and a tree T. This section is a preparation of Section 4. In Section 4, we study the 3-connectivity of the Cartesian product of two connected graphs G and H. Moreover, all the bounds are sharp. Our result could be seen as an extension of Theorem 1.1.

2 Some basic results

We use P_n to denote a path with n vertices. A path P is called a *u-v path*, denoted by $P_{u,v}$, if u and v are the endpoints of P.

Recall that the $Cartesian\ product$ (also called the *square product*) of two graphs G and H, written as $G \,\square\, H$, is the graph with vertex set $V(G) \times V(H)$, in which two vertices (u, v) and (u', v') are adjacent if and only if $u = u'$ and $(v, v') \in E(H)$, or $v = v'$ and $(u, u') \in E(G)$. Clearly, the Cartesian product is commutative, that is, $G \,\square\, H \cong H \,\square\, G$. The edge $(u, v)(u', v')$ is called *one-type edge* if $(u, u') \in E(G)$ and $v = v'$; similarly, the $(u, v)(u', v')$ is called *two-type edge* if $u = u'$ and $(v, v') \in E(H)$.

Let G and H be two graphs with $V(G) = \{u_1, u_2, \ldots, u_n\}$ and $V(H) = \{v_1, v_2, \ldots, v_m\}$, respectively. We use $G(u_j, v_i)$ to denote the subgraph of $G \,\square\, H$ induced by the set $\{(u_j, v_i) \,|\, 1 \leq j \leq n\}$. Similarly, we use $H(u_j, v_i)$ to denote the subgraph of $G \,\square\, H$ induced by the set $\{(u_j, v_i) \,|\, 1 \leq i \leq m\}$. It is easy to see $G(u_{j_1}, v_i) = G(u_{j_2}, v_i)$ for different u_{j_1} and u_{j_2} of G. Thus, we can replace $G(u_j, v_i)$ by $G(v_i)$ for simplicity. Similarly, we can replace $H(u_j, v_i)$ by $H(u_j)$. For any $u, u' \in V(G)$ and $v, v' \in V(G)$, $(u, v),\ (u, v') \in V(H(u))$, $(u', v),\ (u', v') \in V(H(u'))$, $(u, v),\ (u', v) \in V(G(v))$, and $(u, v'),\ (u', v') \in V(G(v))$. We refer to (u, v') and (u', v) as $the\ vertices\ corresponding\ to$ (u, v) in $G(v')\ (= G(u, v')\,)$ and $H(u')\ (= H(u', v)\,)$, respectively. Similarly, we can define the path and tree corresponding to some path and tree, respectively.

In order to show our main results, we need the following well-known theorem.

Theorem 2.1 (Menger's Theorem [1]) *Let G be a k-connected graph, and let x and y be a pair of distinct vertices in G. Then there exist k internally disjoint paths $P_1, P_2, \ldots, P_k$ in G connecting x and y.*

Let G be a connected graph, and $S = \{x_1, x_2, x_3\} \subseteq V(G)$. We first have the following observation about internally disjoint S-trees.

Observation 2.1 *Let G be a connected graph, $S = \{x_1, x_2, x_3\} \subseteq V(G)$, and T be an S-tree. Then there exists a subtree T' of T such that T' is also an S-tree such that $1 \leq d_{T'}(x_i) \leq 2$, $|\{x_i \,|\, d_{T'}(x_i) = 1\}| \geq 2$ and $\{x \,|\, d_{T'}(x) = 1\} \subseteq S$. Moreover, if $|\{x_i \,|\, d_{T'}(x_i) = 1\}| = 3$, then all the vertices of $V(T') \setminus \{x_1, x_2, x_3\}$ have degree 2 except for one vertex, say x with $d_{T'}(x) = 3$; if there exists one vertex of S, say x_1, of degree 2 in T', then T' is an x_2-x_3 path.*

Proof: It is easy to check that this observation holds by deleting vertices and edges of T. □

Remark 2.1 (i) *Since the path between any two distinct vertices is unique in T, the tree T' obtained from T in Observation 2.1 is unique. Such a tree is called a minimal S-tree (or minimal $\{x_1, x_2, x_3\}$-tree).*

(ii) *Let $S = \{x, y, z\} \subseteq V(G)$. Throughout this paper, we can assume that each S-tree is a minimal S-tree.*

Lemma 2.1 *Let G be a graph with $\kappa_3(G) = k \geq 2$, $S = \{x, y, z\} \subseteq V(G)$. Then, the following assertions hold:*

(i) *If $G[S]$ is a clique, then there exist k internally disjoint S-trees $T_1, T_2, \ldots, T_k$, such that $E(T_i) \cap E(G[S]) = \emptyset$ for $1 \leq i \leq k-2$.*

(ii) *If $G[S]$ is not a clique, then there exist k internally disjoint S-trees $T_1, T_2, \ldots, T_k$, such that $E(T_i) \cap E(G[S]) = \emptyset$ for $1 \leq i \leq k-1$.*

Proof: We first prove (i). Clearly, by the definition of S-trees, we know $|\{T_i \mid E(T_i) \cap E(G[S]) \neq \emptyset\}| \leq 3$. Let $\{T_1, T_2, \ldots, T_k\}$ be k internally disjoint S-trees. If $|\{T_i \mid E(T_i) \cap E(G[S]) \neq \emptyset\}| \leq 2$, we are done by exchanging subscript. Thus, suppose $|\{T_i \mid E(T_i) \cap E(G[S]) \neq \emptyset\}| = 3$. Without loss of generality, we assume $E(T_i) \cap E(G[S]) \neq \emptyset$, where $i = k-2, k-1, k$. It is easy to check that T_{k-2}, T_{k-1}, T_k must have the structures as shown in Figures 1 a and c. But, for these two cases, we can obtain T'_{k-2}, T'_{k-1}, T'_k from T_{k-2}, T_{k-1}, T_k, such that $E(T'_{k-2}) \cap \{xy, xz, yz\} = \emptyset$. See Figs. $1b$. and $1d$, where the tree T'_{k-2} is shown by dotted lines. Thus $T_1, T_2, \ldots, T_{k-3}, T'_{k-2}, T'_{k-1}, T'_k$ are our desired S-trees.

The proof of (ii) is similar to that of (i), and thus is omitted. □



Fig. 1: T'_{k-2}, T'_{k-1}, T'_k. An edge is shown by a straight line. The edges (or paths) of a tree are shown by the same type of lines.

Remark 2.2 *Let G be a graph with $\kappa_3(G) = k \geq 2$, $S = \{x, y, z\} \subseteq V(G)$. If $|\{T_i \mid E(T_i) \cap E(G[S]) \neq \emptyset\}| \geq 2$ for any collection $\mathcal{T}$ of k internally disjoint S-trees, then $G[S]$ is a clique. Moreover, $T_{k-1} \cup T_k$ must have the structure as shown in Figure 1e.*

3 The Cartesian product of a connected graph and a path

In this section, we show the following proposition.

Proposition 3.1 *Let G be a graph and P_m be a path with m vertices. The following assertions hold:*
(i) If $\kappa_3(G) = \kappa(G) \geq 1$, then $\kappa_3(G \,\Box\, P_m) \geq \kappa_3(G)$. Moreover, the bound is sharp;
(ii) If $1 \leq \kappa_3(G) < \kappa(G)$, then $\kappa_3(G \,\Box\, P_m) \geq \kappa_3(G) + 1$. Moreover, the bound is sharp.

We shall prove Proposition 3.1 by a series of lemmas. Since the proofs of (i) and (ii) are similar, we only show (ii). Let G be a graph with $V(G) = \{u_1, u_2, \ldots, u_n\}$ such that $1 \leq \kappa_3(G) < \kappa(G)$, $V(P_m) = \{v_1, v_2, \ldots, v_m\}$ such that v_i and v_j are adjacent if and only if $|i - j| = 1$.

Set $\kappa_3(G) = k$ for simplicity. To prove (ii), we need to show that for any $S = \{x, y, z\} \subseteq V(G \,\Box\, H)$, there exist $k + 1$ internally disjoint S-trees. We proceed our proof by the following three lemmas.

Lemma 3.1 *If x, y, z belongs to the same $V(G(v_i))$, $1 \leq i \leq m$, then there exist $k + 1$ internally disjoint S-trees.*

Proof: Without loss of generality, we assume $x, y, z \in V(G(v_1))$. Since $\kappa_3(G) = k$, there exist k internally disjoint S-trees $T_1, T_2, \ldots, T_k$ in $G(v_1)$. We need another S-tree T_{k+1} such that T_{k+1} and T_i are internally disjoint, for $i = 1, 2, \ldots, k$. Let x', y', z' be the vertices corresponding to x, y, z in $G(v_2)$, and T'_1 be the tree corresponding to T_1 in $G(v_2)$. Therefore, tree T_{k+1} obtained from T'_1 by adding three edges xx', yy', zz' is a desired tree. $\Box$

Lemma 3.2 *If only two vertices of $\{x, y, z\}$ belong to some copy $G(v_i)$, then there exist $k + 1$ internally disjoint S-trees.*

Proof: We may assume $x, y \in V(G(v_1)), z \in V(G(v_2))$. In the following argument, we can see that this assumption has no influence on the correctness of our proof. Let x', y' be the vertices corresponding to x, y in $G(v_2)$, z' be the vertex corresponding to z in $G(v_1)$. Consider the following two cases.



Fig. 2: The edges (or paths) of a tree are shown by the same type of lines. The lightest lines stand for edges (or paths) not contained in T_i^*.

Case 1: $z' \notin \{x, y\}$.

Let $S' = \{x, y, z'\}$, and $T_1, T_2, \ldots T_k$ be k internally disjoint S'-trees in $G(v_1)$ such that $|\{T_i \mid E(T_i) \cap E(G(v_1)[S']\} \neq \emptyset|$ is as small as possible. We can assume that $E(T_i) \cap E(G(v_1)[S']) = \emptyset$ for each i, where $1 \leq i \leq k-2$ by Lemma 2.1.

For a tree T_i with $E(T_i) \cap E(G(v_1)[S']) = \emptyset$, let T_i^* be the tree obtained from T_i by adding $z_i z_i'$ and $z_i' z$, and deleting z', where z_i is any one neighbor of z' in T_i, and z_i' is the vertex corresponding to z_i in $G(v_2)$.

If $E(T_k) \cap E(G(v_1)[S']) \neq \emptyset$, say $yz' \in E(T_k) \cap E(G(v_1)[S'])$. Let $T_k^* = T_k + zz'$ and $T_{k+1}^* = T_k' + xx' + yy'$, where T_k' is the tree corresponding to T_k in $G(v_2)$.

If $E(T_{k-1}) \cap E(G(v_1)[S']) \neq \emptyset$ and $E(T_k) \cap E(G(v_1)[S']) \neq \emptyset$. Then $T_{k-1} \cup T_k$ must have one of the structures as shown in Figures 2 a, b and c by Remark 2.2. If T_{k-1} and T_k have the structures as shown in Figure 2a, then we can obtain trees T_{k-1}^*, T_k^* and T_{k+1}^* as shown in Figure 2d. If T_{k-1} and T_k have the structures as shown in Figure 2b, then we can obtain trees T_{k-1}^*, T_k^* and T_{k+1}^* as shown in Figure 2e. If T_{k-1} and T_k have the structures as shown in Figure 2c, then we can obtain trees T_{k-1}^*, T_k^* and T_{k+1}^* similar to those in Figure 2d.

Case 2: $z' \in \{x, y\}$.

Without loss of generality, assume $z' = y$. Since $\kappa(G) > \kappa_3(G) = k$, by Menger's Theorem, there exist at least $k+1$ internally disjoint *x*-*y* paths $P^1, P^2, \ldots, P^{k+1}$ in $G(v_1)$. Assume that y_i is the only neighbor of y in P^i, and that y_i' is the vertex corresponding to y_i in $G(v_2)$. If x and y are nonadjacent in P^i, let T_i be the tree obtained from P^i by adding $y_i y_i'$ and $y_i' z$. If x and y are adjacent in P^i, let T_i be the tree obtained from P^i by adding yz. Since G is a simple graph, there exists at most one path P^i such that x and y are adjacent on P^i. Thus $T_i, 1 \leq i \leq k+1$, are $k+1$ internally disjoint S-trees.

$\square$

Lemma 3.3 *If x, y, z are contained in distinct $G(v_i)$s, then there exist $k+1$ internally disjoint S-trees.*

Proof: We may assume that $x \in V(G(v_1)), y \in V(G(v_2)), z \in V(G(v_3))$. In the following argument, we can see that this assumption has no influence on the correctness of our proof. Let y', z' be the vertices corresponding to y, z in $G(v_1)$, x', z'' be the vertices corresponding to x, z in $G(v_2)$ and x'', y'' be the vertices corresponding to x, y in $G(v_3)$. We consider the following three cases.

Case 1: x, y', z' are distinct vertices in $G(v_1)$

Let $S' = \{x, y', z'\}$, and $T_1, T_2, \ldots T_k$ be k internally disjoint S'-trees in $G(v_1)$ such that $|\{T_i \mid E(T_i) \cap E(G(v_1)[S']) \neq \emptyset\}|$ is as small as possible. We can assume that $E(T_i) \cap E(G(v_1)[S']) = \emptyset$ for each i, where $1 \leq i \leq k-2$ by Lemma 2.1. For each T_i such that $E(T_i) \cap E(G(v_1)[S']) = \emptyset$, we can obtain an S-tree T_i^* from T_i similar to that in Subcase 1.1 of Lemma 3.2.

If $E(T_{k-1}) \cap E(G(v_1)[S']) = \emptyset$ or $E(T_{k-1}) \cap E(G(v_1)[S']) = \emptyset$. Without loss of generality, we assume $E(T_{k-1}) \cap E(G(v_1)[S']) = \emptyset$. Let T_k^* be the tree obtained from T_k by adding edges

$y'y, z'z''$ and $z''z$, T^*_{k+1} be the tree obtained from T''_k by adding $x''x'$, $x'x$ and $y''y$, where T''_k is the tree corresponding to T_k in $G(v_3)$. Thus, T^*_is, $1 \leq i \leq k+1$, are k internally disjoint S-tree.

Otherwise, that is, $E(T_{k-1}) \cap E(G(v_1)[S']) \neq \emptyset$ and $E(T_k) \cap E(G(v_1)[S']) \neq \emptyset$. Then T_{k-1} and T_k must have the structures as shown in Figure 3a, b and c. If T_{k-1} and T_k have the structures as shown in Figure 3a, then we can obtain trees T^*_{k-1}, T^*_k and T^*_{k+1} as shown in Figure 3d. If T_{k-1} and T_k have the structures as shown in Figure 3b, then we can obtain trees T^*_{k-1}, T^*_k and T^*_{k+1} as shown in Figure 3e. If T_{k-1} and T_k have the structures as shown in Figure 3c, then we can obtain trees T^*_{k-1}, T^*_k and T^*_{k+1} as shown in Figure 3f.

Case 2: Two of x, y', z' are the same vertex in $G(v_1)$.

If $y' = z'$, since $\kappa(G) > \kappa_3(G) = k$, by Menger's Theorem, it is easy to construct $k+1$ internally disjoint S-trees. See Figure $3g$. The other cases ($x = y'$ or $x = z'$) can be proved with similar arguments.

Case 3: x, y', z' are the same vertex in $G(v_1)$.

Since $\kappa(G) > \kappa_3(G) = k$, by Menger's Theorem, it is easy to construct $k+1$ internally disjoint S-trees. See Figure 3h.

□

We have the following observation by the argument in the proof of Proposition 3.1.

Observation 3.1 *The $k+1$ internally disjoint S-trees consist of three kinds of edges — the edges of original trees (or paths), the edges corresponding the edges of original trees (or paths) and two-type edges.*

Note that $Q_n \cong P_2 \,\square\, P_2 \,\square \cdots \square\, P_2$, where Q_n is the n-hypercube. We have the following corollary.

Corollary 3.1 *Let Q_n be the n-hypercube with $n \geq 2$. Then $\kappa_3(Q_n) = n-1$.*

Proof: Recall that $\kappa(Q_n) = n$ so that Proposition 3.1 (ii) inductively applies. It is easy to check that $\kappa_3(Q_2) = 1$. Suppose that the result holds for Q_{n-1}, where $n \geq 3$. We have $\kappa_3(Q_n) \geq n-1$ by Proposition 3.1. On the other hand, since Q_n is n-regular, we have $\kappa_3(Q_n) \leq n-1$ by Theorem 1.3. Thus $\kappa_3(Q_n) = n-1$. □

Example 3.1 *Let K_{2n} be the complete graph with vertex set $V(K_{2n}) = \{u_1, u_2, \ldots, u_{2n}\}$, and let G_n be the graph obtained from K_{2n} by adding a new vertex u and edges uu_i, $1 \leq i \leq n$.*

For any $S = \{x, y, z\} \subseteq V(G)$, if $u \notin S$, then there exist k internally disjoint S-trees in G_n by Theorem 1.2. If $u \in S$, without loss of generality, assume $x = u, y = u_1, z = u_2$. Let T_1 be the path u, u_1, u_{k+1}, u_2, T_2 be the path u, u_2, u_{k+2}, u_1, and T_i be the tree obtained from a path u, u_{n+i}, u_1 by adding an edge $u_{n+i}u_2$ for $3 \leq i \leq n$. Clearly, T_i, $1 \leq i \leq n$, are n internally disjoint S-trees. So $\kappa_3(G_n) \geq n$. Since $\delta(G_n) = n$, $\kappa_3(G_n) = n$ by Theorem 1.4. By Proposition 3.1, $\kappa_3(G_n \,\square\, K_2) \geq n$. Since G_n has two adjacent vertices of degree $n+1$, $\kappa_3(G_n) = n$ by Theorem 1.3. Moreover, clearly, $\kappa(G) = n$. Thus $\kappa_3(G \,\square\, K_2) = \kappa_3(G) = n$.

Remark 3.1 *We know that the bounds of (i) and (ii) in Theorem 3.1 are sharp by Example 3.1 and Corollary 3.1.*



Fig. 3: The edges (or paths) of a tree are shown by the same type of lines. The lightest lines stand for edges (or paths) not contained in T_i^*.

Proposition 3.2 *Let G be a connected graph and T be a tree. The following assertions hold:*
(i) *If $\kappa_3(G) = \kappa(G) \geq 1$, then $\kappa_3(G \,\Box\, T) \geq \kappa_3(G)$. Moreover, the bound is sharp;*
(ii) *If $1 \leq \kappa_3(G) < \kappa(G)$, then $\kappa_3(G \,\Box\, T) \geq \kappa_3(G) + 1$. Moreover, the bound is sharp.*

Proof: Since the proofs of (i) and (ii) are similar, we only show (ii). It suffices to show that for any $S = \{x, y, z\} \subseteq V(G \,\Box\, H)$, there exist $k + 1$ internally disjoint S-trees. Set $\kappa_3(G) = k$, $V(G) = \{u_1, u_2, \ldots, u_n\}$, and $V(T) = \{v_1, v_2, \ldots, v_m\}$.

Let $x \in V(G(v_i)), y \in V(G(v_j)), z \in V(G(v_k))$ be three distinct vertices. If there exists a path in T containing v_i, v_j and v_k, then we are done from Proposition 3.1. If i, j and k are not distinct integers, such a path must exist. Thus, suppose that i, j and k are distinct integers, and that there exists no path containing v_i, v_j and v_k. By Observation 2.1, there exists a tree T' in T such that $d_{T'}(v_i) = d_{T'}(v_j) = d_{T'}(v_k) = 1$ and all the vertices of $V(T) \setminus \{v_i, v_j, v_k\}$ have degree 2 except for one vertex, say v_4 with $d_T(v_4) = 3$. Without loss of generality, we set $i = 1, j = 2, k = 3$. Furthermore, we assume $v_iv_4 \in E(T')$, where $1 \leq i \leq 3$. In the following argument, we can see that this assumption has no influence on the correctness of our proof.

Let P be the unique path in T' connecting v_1 and v_2. By Proposition 3.1, we can construct $k + 1$ internally disjoint $\{x, y, z'\}$-trees $T_i,\ 1 \leq k + 1$, in $G \,\Box\, P$, where z' is the vertex corresponding to z in $G(v_4)$. By a similar method of Proposition 3.1, we can construct $k+1$ internally disjoint S-trees in $G \,\Box\, T$ on the basis of these trees. $\Box$

Remark 3.2 *We know that the bounds of (i) and (ii) in Proposition 3.2 are sharp by Example 3.1 and Corollary 3.1.*

Observation 3.2 *The $k + 1$ internally disjoint S-trees consist of three kinds of edges — the edges of original trees (or paths), the edges corresponding the edges of original trees (or paths) and two-type edges.*

4 The Cartesian product of two general graphs

Observation 4.1 *Let G and H be two connected graphs, x, y, z be three distinct vertices in H, and $T_1, T_2, \ldots, T_k$ be k internally disjoint $\{x, y, z\}$-trees in H. Then $G \,\Box\, \bigcup_{i=1}^{k} T_i = \bigcup_{i=1}^{k}(G \,\Box\, T_i)$ has the structure as shown in Figure 4. Moreover, $(G \,\Box\, T_i) \cap (G \,\Box\, T_j) = G(x) \cup G(y) \cup G(z)$ for $i \neq j$. In order to show the structure of $G \,\Box\, \bigcup_{i=1}^{k} T_i$ clearly, we take k copies of $G(y)$, and k copies of $G(z)$. Note that, these k copes of $G(y)$ (resp. $G(z)$) represent the same graph.*

Example 4.1 *Let H be the complete graph of order 4. The structure of $G \,\Box\, (T_1 \cup T_2)$ is shown in Figure 5.*

Now we are ready to prove Theorem 1.5.

Proof of Theorem 1.5: Since the proofs of (i) and (ii) are similar, we only show (ii). Without loss of generality, we set $\kappa_3(G) := k, \kappa_3(H) := \ell$. It suffices to show that for any $S = \{x, y, z\} \subseteq V(G \,\Box\, H)$, there exist $k + \ell$ internally disjoint S-trees. Assume $V(G) = \{u_1, u_2, \ldots, u_n\}$ and $V(T) = \{v_1, v_2, \ldots, v_m\}$.

Let $x \in V(G(v_i)), y \in V(G(v_j)), z \in V(G(v_k))$ be three distinct vertices in $G \,\Box\, H$. We will do only the case that i, j, k are distinct integers. Other two possibilities are similar. Without loss of generality, set $i = 1, j = 2, k = 3$. Since $\kappa_3(H) = \ell$, there exist ℓ internally disjoint $\{v_1, v_2, v_3\}$-trees $T_i,\ 1 \leq i \leq \ell$, in H. We use G_i to denote $G \,\Box\, T_i$. By Observation 4.1, we know that $G \,\Box\, \bigcup_{i=1}^{\ell} T_i = \bigcup_{i=1}^{\ell} G_i$ and



Fig. 4: The structure of $G \,\square\, \bigcup_{i=1}^{k} T_i$.



Fig. 5: The structure of $G \,\square\, (T_1 \cup T_2)$.

$G_i \cap G_j = G(v_1) \cup G(v_2) \cup G(v_3)$ for $i \neq j$. Let y', z' be the vertices corresponding to y, z in $G(v_1)$, respectively.

If x, y', z' are distinct vertices in $G(v_1)$. Since $\kappa_3(G(v_1)) = k$, there exist k internally disjoint $\{x, y', z'\}$-trees T'_j, $1 \leq j \leq k$, in $G(v_1)$. Let $k_0, k_1, \ldots, k_\ell$ be integers such that $0 = k_0 < k_1 < \cdots < k_\ell = k$. Similar to the proofs of Proposition 3.1, we can construct $k_i - k_{i-1} + 1$ internally disjoint S-trees $T_{i,j_i}, 1 \leq j_i \leq k_i - k_{i-1} + 1$, in $(\bigcup_{j=k_{i-1}+1}^{k_i} T'_j) \,\square\, T_i$ for each i, where $1 \leq i \leq \ell$. By Observations 3.1 and 3.2, T_{i,j_i} and T_{r,j_r} are internally disjoint for $i \neq r$. Thus $T_{i,j_i}, 1 \leq i \leq \ell, 1 \leq j_i \leq k_i - k_{i-1} + 1$ are $k + \ell$ internally disjoint S-trees.

If exactly two of x, y', z' are the same vertex in $G(v_i)$. Without loss of generality, assume $y' = z'$. Since $\kappa(G(v_1)) > k$, there exist $k+1$ internally disjoint $x - y'$ paths $P_i,\ 1 \leq i \leq k+1$, in $G(v_1)$ by Menger's Theorem. Note that at most one of them is a path of length 1. Let P_{k+1} be such a path if $xy' \in E(G(v_1))$, and let $k_0, k_1, \ldots, k_\ell$ be integers such that $0 = k_0 < k_1 < \cdots < k_\ell = k+1$. Similar to the proofs of Proposition 3.1, we can construct $k_i - k_{i-1} + 1$ internally disjoint S-trees $T_{i,j_i}, 1 \leq j_i \leq k_i - k_{i-1} + 1$, in $(\bigcup_{j=k_{i-1}+1}^{k_i} P_j) \,\square\, T_i$ for each i, where $1 \leq i \leq \ell - 1$, and $k_\ell - k_{\ell-1}$ internally disjoint S-trees $T_{\ell,j_\ell}, 1 \leq j_i \leq k_\ell - k_{\ell-1}$, in $(\bigcup_{j=k_{\ell-1}+1}^{k_\ell} P_j) \,\square\, T_i$. By Observation 3.1 and 3.2, T_{i,j_i} and T_{r,j_r} are internally disjoint for $i \neq r$. Thus $T_{i,j_i}, 1 \leq i \leq \ell, 1 \leq j_i \leq k_i - k_{i-1} + 1$ are $k + \ell$ internally disjoint S-trees.

If all of x, y', z' are the same vertex in $G(v_i)$. Since $\delta(G(v_1)) \geq \kappa(G(v_1)) > k$, x has k neighbors, say $x_1, x_2, \ldots, x_k$, in $G(v_1))$. Let P_i be the path xx_i, and let $k_0, k_1, \ldots, k_\ell$ be integers such that $0 = k_0 < k_1 < \cdots < k_\ell = k$. Similar to the proofs of Proposition 3.1, we can construct $k_i - k_{i-1} + 1$ internally disjoint S-trees $T_{i,j_i}, 1 \leq j_i \leq k_i - k_{i-1} + 1$, in $(\bigcup_{j=k_{i-1}+1}^{k_i} P_j) \,\square\, T_i$ for each i, where $1 \leq i \leq \ell$. By Observation 3.1 and 3.2, T_{i,j_i} and T_{r,j_r} are internally disjoint for $i \neq r$. Thus $T_{i,j_i}, 1 \leq i \leq \ell, 1 \leq j_i \leq k_i - k_{i-1} + 1$ are $k + \ell$ internally disjoint S-trees.

We now show that bounds of Theorem 1.5 are sharp. For (i), Example 3.1 is a sharp example. Let K_n be a complete graph with n vertices, and P_m be a path with m vertices, where $m \geq 2$. We have $\kappa_3(P_m) = 1$, and $\kappa_3(K_n) = n - 2$ by Theorem 1.2. It is easy to check that $\kappa_3(K_n \,\square\, P_m) = n - 2 + 1 = n - 1$. Thus, $K_n \,\square\, P_m$ is a sharp example for (ii). $\square$

Acknowledgments

We thank anonymous reviewers for their carefully reading of our work and their helpful suggestions.

References

[1] J.A. Bondy, U.S.R. Murty, Graph Theory, GTM 244, Springer, 2008.

[2] G. Chartrand, S.F. Kapoor, L.Lesniak, D.R. Lick, Generalized connectivity in graphs, Bull. Bombay Math. Colloq. 2(1984), 1-6

[3] G. Chartrand, F. Okamoto, P. Zhang, Rainbow trees in graphs and generalized connectivity, Networks 55(4)(2010), 360-367.

[4] W.S. Chiue, B.S. Shieh, On connectivity of the Cartesian product of two graphs. Appl. Math. Comput. 102(1999), 129-137.

[5] W. Imrich, S. Klavžar, Product Graphs Structure and Recongnition, A Wiley-Interscience Publication, 2000.

[6] S. Klavžar, S. Špacapan, On the edge-connectivity of Cartesian product graphs, Asian-Eur. J. Math. 1(2008), 93-98.

[7] S. Li, W. Li, X. Li, The generalized connectivity of complete bipartite graphs, Ars Combin. 104(2012).

[8] S. Li, X. Li, Note on the hardness of generalized connectivity, J. Comb. Optim., doi 10.1007/s10878-011-9399-x.

[9] S. Li, X. Li, W. Zhou, Sharp bounds for the generalized connectivity $\kappa_3(G)$, Discrete Math. 310(2010), 2147-2163.

[10] F. Okamoto, P. Zhang, The tree connectivity of regular complete bipartite graphs, J. Combin. Math. Combin. Comput. 74(2010), 279-293.

[11] G. Sabidussi, Graphs with given group and given graph theoretical properties, Canadian J. Math. 9(1957), 515-525.

[12] S. Špacapan, Connectivity of Cartesian products of graphs. Appl. Math. Lett. 21(2008), 682-685.

[13] N.A. Sherwani, Algorithms for VLSI physical design automation, 3rd Edition, Kluwer Acad. Pub., London, 1999.

[14] H. Whitney, Congruent graphs and the connectivity of graphs, Am. J. Math. 54(1932), 150-168.

[15] I. Wilfried, S. Klavžar, D.F. Rall, Topics in Graph Theory. Graphs and Their Cartesian Product. A K Peters, 2008.

[16] J.M. Xu, C. Yang, Connectivity of Cartesian product graphs. Discrete Math. 306(1)(2006), 159-165