

On the residual solvability of generalized free products of solvable groups

Delaram Kahrobaei^{1,2†} and Stephen Majewicz^{3‡}

¹Doctoral Program in Computer Science, Graduate Center (CUNY), New York, USA

²Mathematics Department, New York City College of Technology (CUNY), New York, USA

³Mathematics Department, Kingsborough Community College (CUNY), New York, USA

received 27th September 2010, revised 29th October 2010, accepted 29th October 2010.

In this paper, we study the residual solvability of the generalized free product of solvable groups.

Keywords: residual solvability, solvable groups, amalgamated products

Dedicated to Laci Babai

1 Introduction and Motivation

In his celebrated paper [4], Hall introduced the residual properties. Let \mathcal{P} be a property of groups, and let G be a group. We say that G is *residually* \mathcal{P} if for any non-identity element $g \in G$, there exists a homomorphism ψ from G to a group having property \mathcal{P} such that $\psi(g) \neq 1$.

We recall the definition of the generalized free products of groups. Let there be given a nonempty set of groups $\{G_\lambda | \lambda \in \Lambda\}$ together with a group H which is isomorphic with a subgroup H_λ of G_λ by means of monomorphism $\phi_\lambda : H \rightarrow G_\lambda$ ($\lambda \in \Lambda$). There is an exceedingly useful object known as the free product of the G_λ 's with the amalgamated subgroup H . Roughly speaking this is the largest group generated by the G_λ 's in which the subgroups H_λ are identified by means of the ϕ_λ . Generically such groups are known as generalized free products (see [8]).

In the literature, residual finiteness has attracted much attention. In [2], for example, Baumslag studies the residual finiteness of generalized free products of nilpotent groups. Another study pursued by M. Sapir [9] and D. Wise [10] focuses on the residual finiteness of one-relator groups.

On the other hand, the exploration of residually solvable groups is in its early stages. One classical work on the residual solvability of positive one-relator groups is due to Baumslag (see [3]). The first author studied doubles of residually solvable groups (see [5]), as well as the residual solvability of the generalized free products of finitely generated nilpotent groups (see [6]). In [1], Arzhantseva, de la Harpe,

[†]Email: dkahrobaei@gc.cuny.edu. Supported by a PSC CUNY Research Foundation of City University of New York and City Tech Foundation.

[‡]Email: smajewicz@kbcc.cuny.edu

and the first author introduce the true prosolvable completion of a group and give several interesting open problems that motivate the importance of residually solvable groups.

In this paper, we generalize some of the results in [6] which carry over when nilpotent groups are replaced by solvable groups.

2 Results

In this section, we state our main results. Their proofs will be presented in Section 4.

We begin by proving that the generalized free product of a nilpotent group and a solvable group is not necessarily perfect.

Theorem 1 *The generalized free product of a nilpotent group and a solvable group amalgamated by a proper subgroup of them is not perfect.*

Next we consider the case where the amalgamated subgroup of the generalized free product of two solvable groups is cyclic. By choosing an appropriate solvable filtration of each factor so that the generator of the amalgamated subgroup does not lie in the n^{th} term of the derived series, it follows that the resulting amalgam is residually solvable.

Theorem 2 *The generalized free product of two solvable groups amalgamating a cyclic subgroup is residually solvable.*

In [6], Kahrobaei proved the generalized free product of two finitely generated solvable groups amalgamated by central subgroups is a solvable extension of a residually solvable group and, hence, is residually solvable. We generalize this result for any finite number of solvable groups.

Theorem 3 *The generalized free product of a finite number of solvable groups, amalgamating a central subgroup in each of the factors, is residually solvable.*

Our next theorem deals with the generalization of doubles (recall that a *double* is the amalgamated product of two groups whose factors are isomorphic and the amalgamated subgroups are identified under the same isomorphism).

Theorem 4 *Let $\{A_i \mid i \in I\}$ be an arbitrary indexed family of isomorphic solvable groups such that $\bigcap_{i \in I} A_i = C$, and let G be the generalized free product of the A_i 's amalgamated by C . Then G is residually solvable.*

Our final result pertains to generalized free products containing a finitely generated torsion-free abelian factor.

Theorem 5 *The generalized free product of a finitely generated torsion-free abelian group and a solvable group is residually solvable.*

Note that Theorem 5 is a special case where the amalgamated subgroup is central in only one of the factors.

3 Background and Preliminary Results

In this section, we present some preliminary material which will be needed for the proofs of our results. We begin with a theorem due to Neumann [7] which plays a crucial role in our work.

Theorem 6 (Neumann) *If K is a subgroup of $G = \{A * B; C\}$, then K is an HNN-extension of a tree product in which the vertex groups are conjugates of subgroups of either A or B and the edge groups are conjugates of subgroups of C . The associated subgroups involved in the HNN-extension are also conjugates of subgroups of C . If $K \cap A = \{1\} = K \cap B$, then K is free. If $K \cap C = \{1\}$, then $K = \prod_{i \in I}^* X_i * F$, where the X_i are conjugates of subgroups of A and B and F is free.*

The next result is used in the proof of Theorem 1. If G is any group, let G_{ab} denote the abelianization of G .

Lemma 7 (Kahrobaei [6]) *Let G be the amalgamated product of A and B with C_A and C_B identified,*

$$G = \{A * B; C_A = C_B\}.$$

Then G_{ab} maps onto

$$D = A_{\text{ab}}/gp(\bar{c}_a \mid c_a \in C_A) \times B_{\text{ab}}/gp(\bar{c}_b^{-1} \mid c_b \in C_B),$$

where \bar{c}_a and \bar{c}_b are the images of c_a and c_b in A_{ab} and B_{ab} , respectively.

Theorem 1 also relies on the next well-known theorem about the Frattini subgroup $\Phi(G)$ of a nilpotent group G . Let $\delta_i G$ denote the i^{th} derived subgroup of a group G .

Theorem 8 (Hirsch) *If G is a nilpotent group, then $\delta_2 G \leq \Phi(G)$.*

The proof of Theorem 3 makes use of the generalized central product of a set of groups. Let $Z(G)$ denote the center of a group G .

Definition 9 *Suppose that*

$$\{A_i = \langle X_i; R_i \rangle \mid i \in I\}$$

is an indexed family of presentations, and let C be a group equipped with monomorphisms

$$\phi_i : C \rightarrow A_i \text{ and } C \leq Z(A_i) \text{ (for all } i \in I\text{)}.$$

The group A defined by the presentation

$$A = \langle \cup X_i; \cup R_i \cup \{c\phi_i c^{-1}\phi_j \mid c \in C, i, j \in I\} \cup \{[x_i, x_j] = 1, i, j \in I\} \rangle,$$

where we assume that the X_i are disjoint, is termed the generalized central product of the A_i amalgamating the central subgroup C . Thus,

$$A = \prod_{i \in I}^{\times} \{A_i; C\}.$$

It is not hard to show that

$$A = \left(\prod_{i \in I}^{\times} A_i \right) / gp(c\phi_i c^{-1} \phi_j \mid i, j \in I, c \in C).$$

By a theorem of von Dyck, there are canonical homomorphisms $\mu_i : A_i \rightarrow A$.

Lemma 10 *Each homomorphism μ_i is a monomorphism.*

We note that the generalized central product of finitely many solvable groups is solvable. For the case of abelian factors, the generalized central product of an arbitrary number of abelian groups is abelian.

A key ingredient for proving Theorem 6 is the following:

Lemma 11 (Kahrobaei [6]) *Suppose $\phi : A \rightarrow B$ is an isomorphism between two groups A and B . Let $C < A$, and let D be the amalgamated product of A and B amalgamating C with $C\phi$:*

$$D = \{A * B; C = C\phi\}$$

There exists a homomorphism ψ from D onto one of the factors with kernel

$$gp(a(a\phi)^{-1} \mid a \in A),$$

and ψ injects into each factor.

4 Proofs of Main Results

Proof of Theorem 1: Let A be a nilpotent group, and let B be a solvable group. Suppose C_A and C_B are proper subgroups of A and B , respectively. We want to show that $G = \{A * B; C_A = C_B\}$ is not perfect.

For any $c_a \in C_A$ and $c_b \in C_B$, let \bar{c}_a and \bar{c}_b denote their images in A_{ab} and B_{ab} , respectively. Set $\bar{C}_A = gp(\bar{c}_a \mid c_a \in C_A)$ and $\bar{C}_B = gp(\bar{c}_b^{-1} \mid c_b \in C_B)$. By Lemma 7, G_{ab} maps onto

$$D = A_{ab}/\bar{C}_A \times B_{ab}/\bar{C}_B.$$

We claim that $A_{ab}/\bar{C}_A \neq \{1\}$. By Theorem 8, $gp(C_A, \delta_2 A)$ is a proper subgroup of A since C_A is. Thus,

$$A_{ab}/\bar{C}_A \simeq (A/\delta_2 A)/(gp(C_A, \delta_2 A)/\delta_2 A) \simeq A/gp(C_A, \delta_2 A) \neq \{1\},$$

proving our claim. It follows that $D \neq \{1\}$ and, hence, $G_{ab} \neq \{1\}$. Therefore, G is not perfect. \square

Proof of Theorem 2: Suppose $A = gp(a_i \mid i \in I)$ and $B = gp(b_j \mid j \in J)$ are solvable groups, and let $a \in A$ and $b \in B$ be non-identity elements. There exists positive integers m and n such that $a \in \delta_m A \setminus \delta_{m+1} A$ and $b \in \delta_n B \setminus \delta_{n+1} B$. Let $G = \{A * B; a = b\}$, and let D be the central product of $A/\delta_{m+1} A$ and $B/\delta_{n+1} B$ amalgamating $a\delta_{m+1} A$ with $b\delta_{n+1} B$:

$$D = \{A/\delta_{m+1} A \times B/\delta_{n+1} B; a\delta_{m+1} A = b\delta_{n+1} B\}.$$

Let $\phi : G \rightarrow D$ be the homomorphism which maps a_i to $(a_i\delta_{m+1} A, \delta_{n+1} B)$ ($i \in I$) and b_j to $(\delta_{m+1} A, b_j\delta_{n+1} B)$ ($j \in J$), and let K be the kernel of ϕ . Then $K \cap C = \{1\}$, where $C = gp(a) = gp(b)$. By Theorem 6, K is a free product of conjugates of subgroups of A and B , and a free group. Thus K is residually solvable and, hence, G is an extension of a residually solvable group by a solvable group. Therefore, G is residually solvable. \square

Proof of Theorem 3: Suppose that $\{A_i \mid i \in I\}$ is a finite indexed family of solvable groups, and let

$$G = \left\{ \prod_{i \in I}^* A_i; C \right\}.$$

Let S be the generalized central product of the A_i :

$$S = \left(\prod_{i \in I}^{\times} A_i \right) / gp(c\phi_i c^{-1}\phi_j \mid i, j \in I, c \in C),$$

where the ϕ_k 's are as in Definition 9. For each $i \in I$, let π_i denote the canonical homomorphism from S to A_i . Since the π_i coincide on C , they can be naturally extended to a homomorphism μ from G into S . Let K be the kernel of μ , and observe that G/K is solvable since S is. Now, μ is one-to-one restricted to each factor; that is,

$$K \cap A_i = \{1\} \text{ for all } i \in I.$$

By Theorem 6, K is free and, consequently, G is a solvable extension of a free group. Therefore, G is residually solvable. \square

Proof of Theorem 4: Let $\{A_i \mid i \in I\}$ be an arbitrary indexed family of isomorphic solvable groups such that $\bigcap_{i \in I} A_i = C$, and let G be the generalized free product of the A_i 's amalgamated by C . Pick an $i \in I$, and let $\phi : G \rightarrow A_i$ be an epimorphism with kernel K . Then K is free because ϕ restricted to each factor is injective by Lemma 11, and

$$A_i \cap K = \{1\} \text{ for all } i \in I.$$

Therefore, G is a solvable extension of a free group and is, thus, residually solvable. \square

Proof of Theorem 5: Let A be a finitely generated torsion-free abelian group, and let B be a solvable group. Suppose

$$G = \{A * B; C\}.$$

Since A is finitely generated torsion-free abelian, C is a direct factor of a subgroup A_1 of A of finite index; that is, $A_1 = C \times H$, where $[A : A_1] = n < \infty$. Let $\{a_i\}$ be a distinct set of coset representatives of A_1 in A ; that is, $A/A_1 = \bigcup_{i=1}^n a_i A_1$. There exists an epimorphism from G onto $\bigcup_{i=1}^n a_i A_1$ with kernel

$$K = gp_G(B, A_1) = \left\{ \prod_{i=1}^n B^{a_i} * A_1; C \right\}.$$

Now let D be the normal closure of finitely many copies of B in K :

$$D = gp_K \left(\bigcup_{i=1}^t B^{a_i} \right) = \left\{ \prod_{i=1}^t B^{a_i}; C \right\}.$$

By Theorem 4, D is residually solvable. Consequently, K is an extension of a residually solvable group by an abelian group. Therefore, G is (residually solvable)-by-abelian and, hence, residually solvable. \square

References

- [1] G. Arzhantseva, P. de la Harpe, D. Kahrobaei, Z. Sunik. The true prosoluble completion of a group: Examples and open problems *Journal of Geometriae Dedicata, Springer Netherlands* **Vol. 124** (2007) No.1, 5–26
- [2] G. Baumslag. On the residual finiteness of generalised free products of nilpotent groups. *Trans. Amer. Math. Soc.* **106** (1963), 193–209.
- [3] G. Baumslag. Positive one-relator groups. *Trans. Amer. Math. Soc.* **156** (1971), 165–183.
- [4] P. Hall. The splitting properties of relatively free groups. *Proc. London Math. Soc.* **4** (1954), 343–356.
- [5] D. Kahrobaei. Doubles of residually solvable groups. *Aspects of Infinite Group Theory, Algebra and Discrete Mathematics, World Scientific* **Vol 1**(2008)
- [6] D. Kahrobaei. Residual Solvability of generalized free products of finitely generated nilpotent groups. *Communications in Algebra, (to appear)*(2010)
- [7] H. Neumann. Generalized free products with amalgamated subgroups. *Amer. J. Math.* **71** (1949), 491–540.
- [8] D. J. S. Robinson. A course in the theory of groups, *Springer-Verlag* (1995)
- [9] M. Sapir. Residual properties of one-relator groups, *arXiv:1001.2829*, (2010)
- [10] D. Wise. The residual finiteness of quasipositive one-relator groups, *J. London Math. Soc.* **66**, (2002), 334–350