# Total domination in $K_5$ - and $K_6$ -covered graphs

## Odile Favaron<sup>1</sup> and H. Karami<sup>2</sup> and S. M. Sheikholeslami<sup>2</sup>

<sup>1</sup>L.R.I., UMR 8623, Université Paris-Sud, F-91405 Orsay, France

received March 31, 2006, revised Dec 20, 2007, accepted Jan 18, 2008.

A graph G is  $K_r$ -covered if each vertex of G is contained in a  $K_r$ -clique. Let  $\gamma_t(G)$  denote the total domination number of G. It has been conjectured that every  $K_r$ -covered graph of order n with no  $K_r$ -component satisfies  $\gamma_t(G) \leq \frac{2n}{r+1}$ . We prove that this conjecture is true for r=5 and 6.

Keywords: total domination, clique cover

### 1 Introduction

Let G be a simple graph with vertex set V(G) and edge set E(G). We use [6] for terminology and notation which are not defined here. The *open neighborhood*  $N_G(v)$  of a vertex  $v \in V(G)$  is the set of all vertices adjacent to v. Its *closed neighborhood* is  $N_G[v] = N_G(v) \cup \{v\}$ . If S is a set of vertices of S, then  $N(S) = \bigcup_{u \in S} N(u)$  and  $N[S] = N(S) \cup S$ . For the sake of simplicity we write S is denoted by S is denoted by S. For an integer S integer S is a multitriangle of order S is the graph consisting of S is denoted by S.

A set  $D \subseteq V$  is a total dominating set if every vertex in V is adjacent to a vertex of D. Obviously every graph without isolated vertices has a total dominating set. The total domination number,  $\gamma_t(G)$ , is the minimum cardinality of a total dominating set. If G has q components  $G_i$ , then  $\gamma_t(G) = \sum_{i=1}^q \gamma_t(G_i)$ . As for the domination number, the determination of the total domination number of a graph is NP-hard and it is interesting to determine good upper bounds on  $\gamma_t(G)$ . Conditions on the density of the graph allow us to lower such bounds. Here we consider the condition that every vertex is contained in a sufficiently large clique.

A  $K_r$ -component of G is a component isomorphic to a clique  $K_r$ . A graph G is  $K_r$ -covered,  $r \geq 2$ , if every vertex of G is contained in a clique  $K_r$ , and minimally  $K_r$ -covered if it is  $K_r$ -covered but G-e is not  $K_r$ -covered for any edge of G. These properties were already considered by Henning and Swart in [5] under the terms "with no  $K_r$ -isolated vertex" or "Property C(1,r)", and "Property C(2,r)", respectively.

1365-8050 © 2008 Discrete Mathematics and Theoretical Computer Science (DMTCS), Nancy, France

<sup>&</sup>lt;sup>2</sup>Department of Mathematics, Azarbaijan University of Tarbiat Moallem, Tabriz, I.R. Iran

In a  $K_r$ -covered graph, a good vertex is a vertex of degree r-1 and a good clique is a clique  $K_r$  containing a good vertex. If z is a good vertex, we denote by  $C_z$  the good clique containing it. The following results, independently proved in [3] and in [4], will be constantly used throughout the paper.

**Theorem A** [3, 4] Every edge of a minimally  $K_r$ -covered graph is contained in a good clique.

In 2004, Cockayne, Favaron and Mynhardt [1] conjectured that every  $K_r$ —covered graph G of order n with no  $K_r$ —component satisfies  $\gamma_t(G) \leq \frac{2n}{r+1}$ . They proved this conjecture for r=3,4. In this paper we prove it for r=5 and 6.

## 2 Proof of the conjecture for r=5 and r=6

The proof uses a particular family  $\mathcal{F}_r$  of minimally  $K_r$ —covered graphs which was already considered in [1, 4]. Recall that the corona of a graph H is obtained from H by adding a pendant edge at each vertex of H and that the middle graph of H is the line graph of the corona of H (see for instance [2]).

**Definition 1**  $\mathcal{F}_r$  is the family of middle graphs of (r-1)-regular graphs.

From this definition  $\mathcal{F}_r$  is the collection of graphs consisting of edge-disjoint cliques of order r, where each such clique contains exactly one vertex of degree r-1 and the remaining r-1 vertices have degree 2(r-1). Let  $\mathcal S$  be the set of these edge-disjoint cliques. Then each vertex of G of degree r-1 belongs to exactly one  $K_r$  in  $\mathcal S$  and each vertex of degree 2(r-1) belongs to exactly two  $K_r$ 's in  $\mathcal S$ .

The following result is proved in [1].

**Theorem B** (See [1]) For  $r \geq 3$ , every graph of order n of  $\mathcal{F}_r$  satisfies  $\gamma_t(G) < \frac{2n}{r+1}$ .

We can now prove the conjecture for r = 5 and r = 6.

**Theorem 1** For r=5 or 6, every  $K_r$ -covered graph G of order n with no  $K_r$ -component satisfies  $\gamma_t(G) \leq \frac{2n}{r+1}$ .

**Proof:** The proof is by induction on n and the first four claims are established for any value of  $r \geq 5$ . The statement is obviously true for the smallest possible order r+1 since then  $\gamma_t(G)=2$ . Suppose the theorem to be true for graphs of order less than n and let G be a  $K_r$ -covered graph with no  $K_r$ -component of order  $n \geq r+2$ . Let F be a minimally  $K_r$ -covered spanning subgraph of G. Since  $\gamma_t(G) \leq \gamma_t(F)$ , it is sufficient to prove  $\gamma_t(F) \leq \frac{2n}{r+1}$ .

If F is not connected but has no  $K_r$ —component, then applying the induction hypothesis to each component of F gives the result.

The case where F is not connected and has some  $K_r$ -components has already been considered in [1]. The second part of the proof of Theorem 6 in [1] establishes by induction on n that a graph having a certain property  $\mathcal{B}(r)$  satisfies  $\gamma_t(G) \leq 2n/(r+1)$ . In the case where a minimal  $K_r$ -covered spanning subgraph F of G has  $K_r$  components, the result is proved without using the induction hypothesis  $\mathcal{B}(r)$ . Hence the same argument holds here. As the proof is rather long, we do not repeat it and refer the reader to [1].

So we suppose now that we are working in a connected minimal  $K_r$ -covered graph F of order  $n \ge r+2$ . Since F is connected, every vertex of F belongs to an edge and thus has a good neighbor by Theorem A. For every pair of adjacent vertices u and v, let

$$P(u,v) = \{x \in N(u,v) \setminus \{u,v\} \mid N(x) \subseteq N(u,v)\}.$$

**Claim 1** If  $|P(u,v)| \ge r - 1$  for some pair of adjacent non-good vertices u and v, then  $\gamma_t(F) \le \frac{2n}{r+1}$ .

**Proof:** Let u' be any good neighbor of u. By Theorem A, the edge uu' is contained in a good clique  $\mathcal C$ . The r-2 neighbors of u' different from u are vertices of  $\mathcal C$  and thus are adjacent to u. So every good neighbor of u, and similarly every good neighbor of v, belongs to P(u,v). Note also that if  $z \in N(u,v) \setminus (P(u,v) \cup \{u,v\})$ , then z has a neighbor  $z_1 \notin N(u,v)$ , and so z belongs to a good clique of the graph  $F' = F[V \setminus (P(u,v) \cup \{u,v\})]$ . Hence F' is  $K_r$ -covered.

Let  $\mathcal{C}_1,\dots,\mathcal{C}_s$  be the  $K_r$ -components of F' if any. Obviously  $(N(w)\setminus V(\mathcal{C}_i))\subseteq P(u,v)\cup \{u,v\}$  for each  $w\in V(\mathcal{C}_i)$ . Since F is connected, each clique  $\mathcal{C}_i$  contains at least one vertex  $w_i$  such that  $(N(w_i)\setminus V(\mathcal{C}_i))\cap (P(u,v)\cup \{u,v\})\neq \emptyset$ . From the definition of P(u,v) we have  $w_i\in N(u,v)$ . Let  $X=P(u,v)\cup \{u,v\}\cup (\cup_{i=1}^s V(\mathcal{C}_i))$ . The graph  $F[V\setminus X]$  is still  $K_r$ -covered and has no  $K_r$ -component. By the induction hypothesis,  $\gamma_t(F[V\setminus X])\leq \frac{2|V\setminus X|}{r+1}$ . Moreover  $\{u,v,w_1,w_2,\cdots,w_s\}$ , or  $\{u,v\}$  if s=0, is a total dominating set of order s+2 of F[X], and |X|=|P(u,v)|+sr+2 with  $s\geq 0$ . Hence if  $|P(u,v)|\geq r-1$ , then  $\gamma_t(F[X])\leq s+2\leq \frac{2|X|}{r+1}$  and we are done.

We suppose henceforth  $|P(u,v)| \le r-2$  for every pair of adjacent non-good vertices of F. Recall that all the good neighbors of u or v belong to P(u,v). If G consists of  $p \ge 2$  cliques  $K_r$  sharing exactly one vertex, then n = p(r-1) + 1 and  $\gamma_t(G) = 2 \le 2n/(r+1)$ . We also suppose in what follows that G has not this structure, which means that every non-good vertex has at least one non-good neighbor.

**Claim 2** Each good clique contains at most r-4 good vertices.

**Proof:** Suppose to the contrary that  $\mathcal{C}$  is a good clique  $(\neq K_r)$  with more than r-4 good vertices. Let  $z_1, z_2, \cdots, z_s$  with  $r-3 \leq s \leq r-1$  be the good vertices and u a non-good vertex of  $\mathcal{C}$ . If u has a non-good neighbor v not in  $\mathcal{C}$ , let  $\mathcal{C}'$  be a good clique containing uv and  $z_1', \cdots, z_t'$   $(1 \leq t \leq r-2)$  the good vertices of  $\mathcal{C}'$ . The vertex v belongs to a second good clique  $\mathcal{C}'' \neq \mathcal{C}$ . Let z'' be a good vertex of  $\mathcal{C}''$ . Then  $\{z_1, \cdots, z_s, z_1', \cdots, z_t', z_t''\}$  is a subset of P(u, v) of order at least  $s+2 \geq r-1$ , a contradiction to  $|P(u, v)| \leq r-2$ . Therefore all the non-good neighbors of u belong to  $\mathcal{C}$ .

Let  $u, u_1, \dots, u_{r-s-1}$  be the non-good vertices of  $\mathcal{C}$  with  $r-s \geq 2$ . Let  $\mathcal{C}'$  ( $\mathcal{C}'_1$  respectively, possibly equal to  $\mathcal{C}'$ ) be a second good clique containing u ( $u_1$ ) and let z' ( $z'_1$ ) be a good vertex of  $\mathcal{C}'$  ( $\mathcal{C}'_1$ ). Then  $\{z_1, \dots, z_s, z', z'_1\} \subseteq P(u, u_1)$ . Since  $|P(u, u_1)| \leq r-2$ , s=r-3,  $z'=z'_1$  and z' is the unique good vertex of the clique  $\mathcal{C}' = \mathcal{C}'_1$ . Among the r-1 non-good vertices of  $\mathcal{C}'$ , at most three are those of  $\mathcal{C}$  and thus at least one, say  $u_2$ , is not in  $\mathcal{C}$ . Let  $\mathcal{C}''$  be a second good clique containing  $u_2$  and z'' a good vertex of  $\mathcal{C}''$ . Then  $\{z_1, \dots, z_{r-3}, z', z''\} \subseteq P(u, u_2)$ , a contradiction which completes the proof.

**Claim 3** No vertex can belong to r-2 good cliques  $K_r$ .

**Proof:** Assume, to the contrary, that a vertex u belongs to r-2 good cliques  $\mathcal{C}_1,\ldots,\mathcal{C}_{r-2}$ , and let  $x_i$  be a good vertex of  $\mathcal{C}_i$  for  $1 \leq i \leq r-2$ . Let w and t be two non-good vertices of  $\mathcal{C}_1 \setminus \{u,x_1\}$ . Since  $\{x_1,\cdots,x_{r-2}\} \subseteq P(u,t)$  and  $|P(u,t)| \leq r-2$ , we have  $w \notin P(u,t)$  and w has a good neighbor w' not in N(u,t) and thus distinct from  $x_1,\cdots,x_{r-2}$ . But then  $\{x_1,\cdots,x_{r-2},w'\} \subseteq P(u,w)$ , which is a contradiction.

**Claim 4** r-3 good cliques  $K_r$  cannot share more than one vertex.

**Proof:** Assume, to the contrary, that  $\mathcal{C}_1, \mathcal{C}_2, \cdots \mathcal{C}_{r-3}$  are r-3 good  $K_r$ 's all containing the (non-good) vertices u and v, and let  $x_i$  be a good vertex of  $\mathcal{C}_i$  for  $1 \leq i \leq r-3$ . From  $|\bigcup_{i=1}^{r-3} V(\mathcal{C}_i)| \geq (r-1) + (r-3) = 2r-4$ , we get  $|\bigcup_{i=1}^{r-3} V(\mathcal{C}_i) \setminus \{x_1, \cdots, x_{r-3}, u, v\}| \geq r-3 \geq 2$  while  $|P(u, v) \setminus \{x_1, \cdots, x_{r-3}\}| \leq 1$ . Therefore at least one vertex z of  $\bigcup_{i=1}^{r-3} V(\mathcal{C}_i) \setminus \{x_1, \cdots, x_{r-3}, u, v\}$  is not in P(u, v). Let z' be a good neighbor of z not in N(u, v). Since  $\{x_1, \cdots, x_{r-3}, z'\} \subseteq P(u, z)$  and  $|P(u, z)| \leq r-2$ , v is not in P(u, z) and thus belongs to a (r-2)th good clique. This contradicts Claim 3.

#### End of the proof of Theorem 1 for r=5

By Claims 2, 3 and 4, each good clique contains exactly one good vertex and four non-good ones, each non-good vertex is contained in exactly two good  $K_5$ 's, and two good  $K_5$ 's intersect in at most one vertex. Therefore the graph F belongs to the family  $\mathcal{F}_5$  described above and thus  $\gamma_t(F) < \frac{2n}{6}$ . This completes the proof for r=5.

#### End of the proof of Theorem 1 for r=6

Henceforth, each good clique is a  $K_6$  containing at most two good vertices.

**Claim 5** Let  $C_1$  and  $C_2$  be two good  $K_6$ 's such that  $|V(C_1) \cap V(C_2)| \geq 2$ . Then

- 1.  $|V(\mathcal{C}) \cap V(\mathcal{C}_i)| \leq 1$  for each other good clique  $\mathcal{C}$  and i = 1, 2;
- 2. the clique  $C_i$  contains exactly one good vertex for i = 1, 2;
- 3. each other good clique C intersecting  $C_1$  or  $C_2$ , contains exactly one good vertex.

#### **Proof:**

(1) Let  $u,v\in V(\mathcal{C}_1)\cap V(\mathcal{C}_2)$  and let  $\mathcal{C}$  be another good clique in G. Assume, to the contrary,  $|V(\mathcal{C}_1)\cap V(\mathcal{C})|\geq 2$ . Let  $x,x_1,x_2$  be good vertices of  $\mathcal{C},\mathcal{C}_1$  and  $\mathcal{C}_2$ , respectively. Suppose  $u\in\mathcal{C}$ . By Claim  $4,v\not\in\mathcal{C}$ . Let  $w\in V(\mathcal{C})\cap V(\mathcal{C}_1)$  and  $w\neq u$ . Since  $\{x,x_1,x_2\}\subseteq P(u,v)$  and  $|P(u,v)|\leq 4$ , at least one vertex t of  $V(\mathcal{C}_1)\setminus\{u,v,x_1,w\}$  is not in P(u,v). Let t' be a good neighbor of t not in N(u,v). Now we have  $\{x,x_1,x_2,t'\}\subseteq P(u,t)$ . Since  $|P(u,t)|\leq 4$ , w is not in P(u,t) and has a good neighbor w' not in N(u,t). Then  $\{x,x_1,x_2,w'\}\subseteq P(u,w)$ . Thus v is not in P(u,w) and has a good neighbor v' not in N(u,w). This implies  $\{x,x_1,x_2,w',v'\}\subseteq P(v,w)$  which is a contradiction. Thus  $u,v\not\in V(\mathcal{C})$ . Let  $w_1,w_2\in V(\mathcal{C})\cap V(\mathcal{C}_1)$ . Since  $\{x,x_1,x_2\}\subseteq P(u,w_1)$  and  $|P(u,w_1)|\leq 4$ ,  $v\not\in P(u,w_1)$  or  $w_2\not\in P(u,w_1)$ .

First let  $v \notin P(u, w_1)$ . Let v' be a good neighbor of v not in  $N(u, w_1)$ . Now we have  $\{x, x_1, x_2, v'\} \subseteq P(v, w_1)$ . Since  $|P(v, w_1)| \le 4$ ,  $w_2$  is not in  $P(v, w_1)$  and has a good neighbor  $w_2'$  not in  $N(v, w_1)$ . Now we have  $\{x, x_1, x_2, v', w_2'\} \subseteq P(v, w_2)$  which is a contradiction.

Now let  $w_2 \notin P(u, w_1)$ . Let  $w_2'$  be a good neighbor of  $w_2$  not in  $N(u, w_1)$ . Now we have  $\{x, x_1, x_2, w_2'\} \subseteq P(u, w_2)$ . Since  $|P(u, w_2)| \le 4$ , v is not in  $P(u, w_2)$  and has a good neighbor v' not in  $N(u, w_2)$ . This implies that  $|P(v, w_2)| \ge 5$  which is a contradiction.

(2) Suppose  $\mathcal{C}_1$  contains two good vertices  $x_1$  and  $x_1'$ . Since  $|P(u,v)| \leq 4$ ,  $V(\mathcal{C}_1 \cup \mathcal{C}_2) \setminus \{u,v\}$  has a non-good vertex, say w, not in P(u,v). Let w' be a good neighbor of w not in N(u,v). Then  $\{x_1,x_1',x_2,w'\}\subseteq P(u,w)$ . Hence v is not in P(u,w) and has a good neighbor  $v'\notin N(u,w)$ , which implies  $|P(v,w)|\geq 5$ , a contradiction.

(3) Suppose  $\mathcal C$  contains two good vertices y and y'. If  $\mathcal C$  intersects  $\mathcal C_1 \cup \mathcal C_2$  in u, then  $\{x_1, x_2, y, y'\} \subseteq P(u, v)$  and there exists a vertex t of  $V(\mathcal C_1 \cup \mathcal C_2) \setminus \{u, v\}$  with a good neighbor t' not in N(u, v). Then  $\{x_1, x_2, y, y', t'\} \subseteq P(u, t)$ , a contradiction. If  $\mathcal C$  intersects  $\mathcal C_1 \cup \mathcal C_2$  in w different from u and v, then, since  $\{x_1, x_2, y, y'\} \subseteq P(u, w)$ , v has a good neighbor v' not belonging to N(u, w). Hence  $\{x_1, x_2, y, y', v'\} \subseteq P(v, w)$ , a contradiction.

**Claim 6** Let three good cliques  $C_1$ ,  $C_2$  and  $C_3$  share one vertex u. Then

- 1.  $|V(\mathcal{C}) \cap V(\mathcal{C}_i)| \leq 1$  for each other good clique  $\mathcal{C}$  and i = 1, 2, 3;
- 2. for i = 1, 2, 3, each non-good vertex of  $C_i \setminus \{u\}$  belongs to exactly two good cliques;
- 3. for i = 1, 2, 3, each clique  $C_i$  and each good clique C intersecting one of the  $C_i$ 's contains exactly one good vertex.

#### **Proof:**

- (1) Suppose that  $\mathcal{C}$  is a good clique such that  $|V(\mathcal{C}) \cap V(\mathcal{C}_1)| \geq 2$ . By claim 3,  $u \notin V(\mathcal{C})$ . Let  $v, w \in V(\mathcal{C}) \cap V(\mathcal{C}_1)$ . Let  $x, x_1, x_2, x_3$  be good vertices of  $\mathcal{C}, \mathcal{C}_1, \mathcal{C}_2$  and  $\mathcal{C}_3$ , respectively. We have  $\{x, x_1, x_2, x_3\} \subseteq P(u, v)$  and so w is not in P(u, v) and has a good neighbor w' not in N(u, v). Then we have  $|P(u, w)| \geq 5$  which is a contradiction.
- (2) If a vertex  $v \neq u$  of some  $C_i$  belongs to two other good cliques, let v' and v'' two good neighbors of v respectively belonging to these two cliques. Then  $\{x_1, x_2, x_3, v', v''\} \subseteq P(u, v)$ , a contradiction.
- (3) If, say,  $C_1$  has a second good vertex  $x_1'$ , then  $C_2$  and  $C_3$  have one good vertex each, for otherwise  $|P(u,v)| \geq 5$  for any neighbor v of u. Hence there exists at least one non-good vertex v belonging to exactly one of the  $C_i$ 's. This vertex v has a good neighbor  $v' \notin \{x_1, x_1', x_2, x_3\}$  and  $|P(u,v)| \geq 5$ , a contradiction. If a good clique C intersecting one of the  $C_i$ 's in one vertex v (necessarily different from u) contains two good neighbors x and x', then  $\{x_1, x_2, x_3, x, x'\} \subseteq P(u, v)$ , a contradiction.

**Claim 7** Let C be a good clique containing two good vertices  $z_1, z_2$ . Then

- 1. each good clique intersects C in at most one vertex;
- 2. each non-good vertex of C belongs to exactly two good cliques;
- 3. if  $\mathcal{C}'$  is a good clique intersecting  $\mathcal{C}$  in u, then  $\mathcal{C}'$  contains exactly one good vertex, each non-good vertex of  $\mathcal{C}'$  belongs to exactly two good cliques,  $|V(\mathcal{C}') \cap V(\mathcal{C}_1)| \leq 1$  for each good clique  $\mathcal{C}_1$  and if  $|V(\mathcal{C}') \cap V(\mathcal{C}_1)| = 1$ , then  $\mathcal{C}_1$  contains exactly one good vertex.

**Proof:** (1) and (2) are consequences of Claim 5 (2) and 6 (3).

(3) Let u' be a good vertex of  $\mathcal{C}'$ , w be a non-good vertex in  $V(\mathcal{C}') \setminus \{u\}$  and w' a good neighbor of w not in N(u). If w has another good neighbor w'', which can be either a second good vertex of  $\mathcal{C}'$  or of a second clique  $\mathcal{C}_1$  containing w, or a good vertex of a third good clique containing w, then  $\{z_1, z_2, u', w', w''\} \subseteq P(u, w)$ , a contradiction. If a good clique  $\mathcal{C}_1$  intersects  $\mathcal{C}'$  in v and w (both different from u by (2)), then  $v \notin P(u, w)$  since  $\{z_1, z_2, u', w'\} \subseteq P(u, w)$ . Therefore v has another good neighbor  $v' \notin N(u, w)$  and  $\{z_1, z_2, u', w', v'\} \subseteq P(u, v)$ , a contradiction.

Let  $V_i = \{u \in V(F) \mid u \text{ belongs to exactly } i \text{ good cliques}\}, i = 1, 2, 3$ . By Claim 2,  $V_1, V_2, V_3$  partition V(F). Obviously  $V_1$  consists of all good vertices of F. Let t be the number of good cliques that contain two good vertices. Counting the number of edges of F with one endpoint in  $V_1$  and another in  $V_2 \cup V_3$ , implies by Claim 7 that

$$5|V_1| - 2t = 2|V_2| + 4t + 3|V_3|.$$

On the other hand, we have

$$2n = 2(|V_1| + |V_2| + |V_3|).$$

It follows from the last two equations

$$|V_1| = \frac{2n}{7} + \frac{|V_3| + 6t}{7} \ . \tag{1}$$

The following claim gives the structure of the subgraph induced by  $V_3$  in F.

**Claim 8**  $F[V_3]$  is a disjoint union of s cliques with  $s \ge |V_3|/5$ .

**Proof:** Let u and v be two adjacent vertices in  $V_3$ . If the edge uv belongs to only one good clique  $\mathcal{C}_z$ , let  $\mathcal{C}_{u'}$  and  $\mathcal{C}_{u''}$  (respectively  $\mathcal{C}_{v'}$  and  $\mathcal{C}_{v''}$ ) be the other two good cliques containing u (respectively v). Then  $\{u', u'', v', v'', z\}$  is a set of five vertices contained in P(u, v), a contradiction. Therefore every edge joining two vertices in  $V_3$  is contained in exactly (by Claim 5) two good cliques. Let now uvw be a path of  $F[V_3]$ . Among the three good cliques containing v, two contain uv and two contain vw. Hence one of them, say  $\mathcal{C}_z$ , contains  $\{u, v, w\}$  and u and w are adjacent. Moreover, the second good cliques respectively containing uv and vw are the same by Claim 5. This implies that  $\{u, v, w\}$  is contained in exactly two good cliques  $\mathcal{C}_z$  and  $\mathcal{C}_{z'}$ . The preceding arguments show that  $F[V_3]$  is a disjoint union of s cliques  $Q_i$ . Each  $Q_i$  is a part of the intersection of two good cliques  $\mathcal{C}_z$  and  $\mathcal{C}_{z'}$ , thus implying  $|Q_i| \leq 5$ , and each vertex u of  $Q_i$  belongs to a third clique intersecting  $\mathcal{C}_z$  and  $\mathcal{C}_{z'}$  exactly in u. Finally, since  $|Q_i| \leq 5$ ,  $s \geq |V_3|/5$ .

We define now the graph  $F^*$  with vertex set  $\{z \in V(F) \mid z \text{ is a good vertex in } F\}$  and two vertices of  $F^*$  are adjacent if and only if they belong to the same clique or their corresponding good cliques have a common vertex. Since F is connected and each edge of F belongs to a good clique, the graph  $F^*$  is connected.

Three good vertices  $z_1, z_2, z_3$  form a triangle in  $F^*$  if and only if

- 1. the good cliques  $C_{z_1}$ ,  $C_{z_2}$  and  $C_{z_3}$  are different and share one vertex,
- 2. or, say,  $C_{z_1} = C_{z_2}$  and  $C_{z_1} \cap C_{z_3} \neq \emptyset$ ,

3. or the three cliques are pairwise intersecting but  $C_{z_1} \cap C_{z_2} \cap C_{z_3} = \emptyset$ .

We only consider the triangles of the first two types and call them respectively 1-triangles and 2-triangles.

A 1-triangle of  $F^*$  comes from a vertex of  $V_3$ . From Claim 6 (2), if two 1-triangles  $z_1z_2z_3$  and  $z_1'z_2'z_3'$  are not disjoint, then they share one edge, say,  $z_1=z_1'$  and  $z_2=z_2'$ . From Claim 8,  $|V(\mathcal{C}_{z_1})\cap V(\mathcal{C}_{z_2})|\geq 2$  and each good clique  $\mathcal{C}_{z_3}$  and  $\mathcal{C}_{z_3'}$  shares one vertex with  $\mathcal{C}_{z_1}$  and  $\mathcal{C}_{z_2}$ . Since  $|V(\mathcal{C}_{z_1})\cap V(\mathcal{C}_{z_2})|\leq 5$ , at most five 1-triangles share a common edge. Hence the 1-triangles of  $F^*$  form multitriangles MT $_i$  of respective order  $p_i$  with  $1\leq p_i\leq 7$ . We call them multitriangles of type 1 and we associate to each of them the clique  $Q_i\subseteq V(\mathcal{C}_{z_1})\cap V(\mathcal{C}_{z_2})$  of order  $p_i-2\leq 5$  as described in Claim 8. Therefore there are  $s\geq |V_3|/5$  multitriangles of type 1 and all of them are disjoint.

A 2-triangle of  $F^*$  comes from a good clique  $\mathcal{C}$  of F with two good vertices  $z_1$  and  $z_2$ . By Claim 7, the four other vertices of  $\mathcal{C}$  belong to exactly one other good clique and these four good cliques are different. Hence the edge  $z_1z_2$  belongs to exactly four 2-triangles forming a multitriangle of order  $p_i=6$ , called multitriangle of type 2. To each multitriangle MT<sub>i</sub> of type 2 we associate the clique  $Q_i$  of order  $p_i-2=4$  of F formed by the non-good vertices of  $\mathcal{C}$ . There are t multitriangles of type 2, the number of good cliques with two good vertices. By Claim 7, they are pairwise disjoint and disjoint from the multitriangles of type 1.

Let  $F^{**}$  be a spanning subgraph of  $F^*$  containing all the edges of the multitriangles but no other cycle

(the edges of  $F^{**}$  not in multitriangles form a spanning tree of the graph of order  $|V_1| - \sum_{i=1}^{s+i} (p_i - 1)$ 

obtained from  $F^*$  by contracting each multitriangle into one vertex). We form a subset D of vertices of F as follows. For each multitriangle  $\operatorname{MT}_i$  of order  $p_i$ ,  $0 \le i \le s+t$ , put in D the  $p_i-2$  vertices of its associated clique  $Q_i$ . For each edge  $z_iz_j$  of  $F^{**}$  not in a multitriangle, put in D one vertex of  $\mathcal{C}_{z_i} \cap \mathcal{C}_{z_j}$ . The induced subgraph F[D] is connected since  $F^{**}$  is connected and can be seen as the graph representative of the 1- and 2-triangles and of the cutting edges of  $F^{**}$ . The set D contains a vertex in

each good clique and thus dominates F. Hence  $\gamma_t(F) \leq |D|$ . Since  $F^{**}$  contains  $|V_1| - \sum_{i=1}^{s+t} (p_i - 1) - 1$  cutting edges,

$$|D| = \sum_{i=1}^{s+t} (p_i - 2) + |V_1| - \sum_{i=1}^{s+t} (p_i - 1) - 1 = |V_1| - s - t - 1$$

with  $s \ge |V_3|/5$ . By (1) we get

$$|D| \le \frac{2n}{7} + \frac{|V_3|}{7} + \frac{6t}{7} - \frac{|V_3|}{5} - t - 1 < \frac{2n}{7}$$
.

This completes the proof of Theorem 1 for r = 6.

## References

- [1] E. J. Cockayne, O. Favaron, and C. M. Mynhardt, *Total domination in*  $K_r$ -covered graphs , Ars Combin. **71** (2004), 289-303.
- [2] E. J. Cockayne, S. T. Hedetniemi, and D. J. Miller, *Properties of hereditary hypergraphs and middle graphs*, Canad. Math. Bull. **21(4)** (1978), 461-468.
- [3] R. C. Entringer, W. Goddard and M. A. Henning, *A note on cliques and independent sets*, J. Graph Theory **24** (1997), 21-23.
- [4] O. Favaron, H. Li and M. D. Plummer, *Some results on*  $K_r$ -covered graphs, Utilitas Math. **54** (1998), 33-44.
- [5] M. A. Henning and H. C. Swart, *Bounds on a generalized domination parameter*, Questiones Math. **13** (1990), 237-253.
- [6] D.B. West, *Introduction to Graph Theory*, Prentice-Hall, Inc, 2000.