A q-analog of Ljunggren's binomial congruence

Armin Straub[1†]

[1] *Tulane University, New Orleans, LA, USA*

Abstract. We prove a q-analog of a classical binomial congruence due to Ljunggren which states that

$$\binom{ap}{bp} \equiv \binom{a}{b}$$

modulo p^3 for primes $p \geqslant 5$. This congruence subsumes and builds on earlier congruences by Babbage, Wolstenholme and Glaisher for which we recall existing q-analogs. Our congruence generalizes an earlier result of Clark.

Résumé. Nous démontrons un q-analogue d'une congruence binomiale classique de Ljunggren qui stipule:

$$\binom{ap}{bp} \equiv \binom{a}{b}$$

modulo p^3 pour p premier tel que $p \geqslant 5$. Cette congruence s'inspire d'une précédente congruence prouvée par Babbage, Wolstenholme et Glaisher pour laquelle nous présentons les q-analogues existantes. Notre congruence généralise un précédent résultat de Clark.

Keywords: q-analogs, binomial coefficients, binomial congruence

---

1 Introduction and notation

Recently, q-analogs of classical congruences have been studied by several authors including (Cla95), (And99), (SP07), (Pan07), (CP08), (Dil08). Here, we consider the classical congruence

$$\binom{ap}{bp} \equiv \binom{a}{b} \mod p^3 \tag{1}$$

which holds true for primes $p \geqslant 5$. This also appears as Problem 1.6 (d) in (Sta97). Congruence (1) was proved in 1952 by Ljunggren, see (Gra97), and subsequently generalized by Jacobsthal, see Remark 6.

---

†Partially supported by grant NSF-DMS 0713836. astraub@tulane.edu

Let $[n]_q := 1 + q + \ldots q^{n-1}$, $[n]_q! := [n]_q[n-1]_q \cdots [1]_q$ and

$$\binom{n}{k}_q := \frac{[n]_q!}{[k]_q![n-k]_q!}$$

denote the usual q-analogs of numbers, factorials and binomial coefficients respectively. Observe that $[n]_1 = n$ so that in the case $q = 1$ we recover the usual factorials and binomial coefficients as well. Also, recall that the q-binomial coefficients are polynomials in q with nonnegative integer coefficients. An introduction to these q-analogs can be found in (Sta97).

We establish the following q-analog of (1):

Theorem 1 *For primes $p \geqslant 5$ and nonnegative integers a, b,*

$$\binom{ap}{bp}_q \equiv \binom{a}{b}_{q^{p^2}} - \binom{a}{b+1}\binom{b+1}{2}\frac{p^2-1}{12}(q^p-1)^2 \mod [p]_q^3. \tag{2}$$

The congruence (2) and similar ones to follow are to be understood over the ring of polynomials in q with integer coefficients. We remark that $p^2 - 1$ is divisible by 12 for all primes $p \geqslant 5$.

Observe that (2) is indeed a q-analog of (1): as $q \to 1$ we recover (1).

Example 2 Choosing $p = 13$, $a = 2$, and $b = 1$, we have

$$\binom{26}{13}_q = 1 + q^{169} - 14(q^{13}-1)^2 + (1 + q + \ldots + q^{12})^3 f(q)$$

where $f(q) = 14 - 41q + 41q^2 - \ldots + q^{132}$ is an irreducible polynomial with integer coefficients. Upon setting $q = 1$, we obtain $\binom{26}{13} \equiv 2$ modulo 13^3.

Since our treatment very much parallels the classical case, we give a brief history of the congruence (1) in the next section before turning to the proof of Theorem 1.

2 A bit of history

A classical result of Wilson states that $(n-1)! + 1$ is divisible by n if and only if n is a prime number. "In attempting to discover some analogous expression which should be divisible by n^2, whenever n is a prime, but not divisible if n is a composite number", (Bab19), Babbage is led to the congruence

$$\binom{2p-1}{p-1} \equiv 1 \mod p^2 \tag{3}$$

for primes $p \geqslant 3$. In 1862 Wolstenholme, (Wol62), discovered (3) to hold modulo p^3, "for several cases, in testing numerically a result of certain investigations, and after some trouble succeeded in proving it to hold universally" for $p \geqslant 5$. To this end, he proves the fractional congruences

$$\sum_{i=1}^{p-1} \frac{1}{i} \equiv 0 \mod p^2, \tag{4}$$

$$\sum_{i=1}^{p-1} \frac{1}{i^2} \equiv 0 \mod p \tag{5}$$

for primes $p \geqslant 5$. Using (4) and (5) he then extends Babbage's congruence (3) to hold modulo p^3:

$$\binom{2p-1}{p-1} \equiv 1 \mod p^3 \tag{6}$$

for all primes $p \geqslant 5$. Note that (6) can be rewritten as $\binom{2p}{p} \equiv 2$ modulo p^3. The further generalization of (6) to (1), according to (Gra97), was found by Ljunggren in 1952. The case $b = 1$ of (1) was obtained by Glaisher, (Gla00), in 1900.

In fact, Wolstenholme's congruence (6) is central to the further generalization (1). This is just as true when considering the q-analogs of these congruences as we will see here in Lemma 5.

A q-analog of the congruence of Babbage has been found by Clark (Cla95) who proved that

$$\binom{ap}{bp}_q \equiv \binom{a}{b}_{q^{p^2}} \mod [p]_q^2. \tag{7}$$

We generalize this congruence to obtain the q-analog (2) of Ljunggren's congruence (1). A result similar to (7) has also been given by Andrews in (And99).

Our proof of the q-analog proceeds very closely to the history just outlined. Besides the q-analog (7) of Babbage's congruence (3) we will employ q-analogs of Wolstenholme's harmonic congruences (4) and (5) which were recently supplied by Shi and Pan, (SP07):

Theorem 3 *For primes* $p \geqslant 5$,

$$\sum_{i=1}^{p-1} \frac{1}{[i]_q} \equiv -\frac{p-1}{2}(q-1) + \frac{p^2-1}{24}(q-1)^2[p]_q \mod [p]_q^2 \tag{8}$$

as well as

$$\sum_{i=1}^{p-1} \frac{1}{[i]_q^2} \equiv -\frac{(p-1)(p-5)}{12}(q-1)^2 \mod [p]_q. \tag{9}$$

This generalizes an earlier result (And99) of Andrews.

3 A q-analog of Ljunggren's congruence

In the classical case, the typical proof of Ljunggren's congruence (1) starts with the Chu-Vandermonde identity which has the following well-known q-analog:

Theorem 4

$$\binom{m+n}{k}_q = \sum_j \binom{m}{j}_q \binom{n}{k-j}_q q^{j(n-k+j)}.$$

We are now in a position to prove the q-analog of (1).

Proof of Theorem 1: As in (Cla95) we start with the identity

$$\binom{ap}{bp}_q = \sum_{c_1+\ldots+c_a=bp} \binom{p}{c_1}_q \binom{p}{c_2}_q \cdots \binom{p}{c_a}_q q^{p\sum_{1\leqslant i\leqslant a}(i-1)c_i - \sum_{1\leqslant i<j\leqslant a} c_i c_j} \tag{10}$$

which follows inductively from the q-analog of the Chu-Vandermonde identity given in Theorem 4. The summands which are not divisible by $[p]_q^2$ correspond to the c_i taking only the values 0 and p. Since each such summand is determined by the indices $1 \leqslant j_1 < j_2 < \ldots < j_b \leqslant a$ for which $c_i = p$, the total contribution of these terms is

$$\sum_{1 \leqslant j_1 < \ldots < j_b \leqslant a} q^{p^2 \sum_{k=1}^{b} (j_k - 1) - p^2 \binom{b}{2}} = \sum_{0 \leqslant i_1 \leqslant \ldots \leqslant i_b \leqslant a-b} q^{p^2 \sum_{k=1}^{b} i_k} = \binom{a}{b}_{q^{p^2}}.$$

This completes the proof of (7) given in (Cla95).

To obtain (2) we now consider those summands in (10) which are divisible by $[p]_q^2$ but not divisible by $[p]_q^3$. These correspond to all but two of the c_i taking values 0 or p. More precisely, such a summand is determined by indices $1 \leqslant j_1 < j_2 < \ldots < j_b < j_{b+1} \leqslant a$, two subindices $1 \leqslant k < \ell \leqslant b+1$, and $1 \leqslant d \leqslant p-1$ such that

$$c_i = \begin{cases} d \text{ for } i = j_k, \\ p - d \text{ for } i = j_\ell, \\ p \text{ for } i \in \{j_1, \ldots, j_{b+1}\} \backslash \{j_k, j_\ell\}, \\ 0 \text{ for } i \notin \{j_1, \ldots, j_{b+1}\}. \end{cases}$$

For each fixed choice of the j_i and k, ℓ the contribution of the corresponding summands is

$$\sum_{d=1}^{p-1} \binom{p}{d}_q \binom{p}{p-d}_q q^{p \sum_{1 \leqslant i \leqslant a} (i-1) c_i - \sum_{1 \leqslant i < j \leqslant a} c_i c_j}$$

which, using that $q^p \equiv 1$ modulo $[p]_q$, reduces modulo $[p]_q^3$ to

$$\sum_{d=1}^{p-1} \binom{p}{d}_q \binom{p}{p-d}_q q^{d^2} = \binom{2p}{p}_q - [2]_{q^{p^2}}.$$

We conclude that

$$\binom{ap}{bp}_q \equiv \binom{a}{b}_{q^{p^2}} + \binom{a}{b+1} \binom{b+1}{2} \left(\binom{2p}{p}_q - [2]_{q^{p^2}} \right) \mod [p]_q^3. \tag{11}$$

The general result therefore follows from the special case $a = 2$, $b = 1$ which is separately proved next. □

4 A q-analog of Wolstenholme's congruence

We have thus shown that, as in the classical case, the congruence (2) can be reduced, via (11), to the case $a = 2$, $b = 1$. The next result therefore is a q-analog of Wolstenholme's congruence (6).

Lemma 5 *For primes $p \geqslant 5$,*

$$\binom{2p}{p}_q \equiv [2]_{q^{p^2}} - \frac{p^2 - 1}{12} (q^p - 1)^2 \mod [p]_q^3.$$

Proof: Using that $[an]_q = [a]_{q^n}\,[n]_q$ and $[n+m]_q = [n]_q + q^n\,[m]_q$ we compute

$$\binom{2p}{p}_q = \frac{[2p]_q\,[2p-1]_q\cdots[p+1]_q}{[p]_q\,[p-1]_q\cdots[1]_q} = \frac{[2]_{q^p}}{[p-1]_q!}\prod_{k=1}^{p-1}\left([p]_q + q^p\,[p-k]_q\right)$$

which modulo $[p]_q^3$ reduces to (note that $[p-1]_q!$ is relatively prime to $[p]_q^3$)

$$[2]_{q^p}\left(q^{(p-1)p} + q^{(p-2)p}\sum_{1\leqslant i\leqslant p-1}\frac{[p]_q}{[i]_q} + q^{(p-3)p}\sum_{1\leqslant i<j\leqslant p-1}\frac{[p]_q\,[p]_q}{[i]_q\,[j]_q}\right). \tag{12}$$

Combining the results (8) and (9) of Shi and Pan, (SP07), given in Theorem 3, we deduce that for primes $p\geqslant 5$,

$$\sum_{1\leqslant i<j\leqslant p-1}\frac{1}{[i]_q\,[j]_q} \equiv \frac{(p-1)(p-2)}{6}(q-1)^2 \mod [p]_q. \tag{13}$$

Together with (8) this allows us to rewrite (12) modulo $[p]_q^3$ as

$$[2]_{q^p}\left(q^{(p-1)p} + q^{(p-2)p}\left(-\frac{p-1}{2}(q^p-1) + \frac{p^2-1}{24}(q^p-1)^2\right) + \right.$$
$$\left. + q^{(p-3)p}\frac{(p-1)(p-2)}{6}(q^p-1)^2\right).$$

Using the binomial expansion

$$q^{mp} = ((q^p-1)+1)^m = \sum_k \binom{m}{k}(q^p-1)^k$$

to reduce the terms q^{mp} as well as $[2]_{q^p} = 1+q^p$ modulo the appropriate power of $[p]_q$ we obtain

$$\binom{2p}{p}_q \equiv 2 + p(q^p-1) + \frac{(p-1)(5p-1)}{12}(q^p-1)^2 \mod [p]_q^3.$$

Since

$$[2]_{q^{p^2}} \equiv 2 + p(q^p-1) + \frac{(p-1)p}{2}(q^p-1)^2 \mod [p]_q^3$$

the result follows. □

Remark 6 Jacobsthal, see (Gra97), generalized the congruence (1) to hold modulo p^{3+r} where r is the p-adic valuation of

$$ab(a-b)\binom{a}{b} = 2a\binom{a}{b+1}\binom{b+1}{2}.$$

It would be interesting to see if this generalization has a nice analog in the q-world.

Acknowledgements

Most parts of this paper have been written during a visit of the author at Grinnell College. The author wishes to thank Marc Chamberland for his encouraging and helpful support. Partial support of grant NSF-DMS 0713836 is also thankfully acknowledged.

References

[And99] George E. Andrews. q-analogs of the binomial coefficient congruences of Babbage, Wolstenholme and Glaisher. *Discrete Math.*, 204(1):15–25, 1999.

[Bab19] Charles Babbage. Demonstration of a theorem relating to prime numbers. *Edinburgh Philosophical J.*, 1:46–49, 1819.

[Cla95] W. Edwin Clark. q-analogue of a binomial coefficient congruence. *Internat. J. Math. and Math. Sci.*, 18(1):197–200, 1995.

[CP08] Robin Chapman and Hao Pan. q-analogues of Wilson's theorem. *Int. J. Number Theory*, 4(4):539–547, 2008.

[Dil08] Karl Dilcher. Determinant expressions for q-harmonic congruences and degenerate Bernoulli numbers. *Electron. J. Combin.*, 15(1), 2008.

[Gla00] James W. L. Glaisher. Residues of binomial-theorem coefficients with respect to p^3. *Quart. J. Math., Oxford Ser.*, 31:110–124, 1900.

[Gra97] Andrew Granville. Arithmetic properties of binomial coefficients I: Binomial coefficients modulo prime powers. *CMS Conf. Proc.*, 20:253–275, 1997.

[Pan07] Hao Pan. A q-analogue of Lehmer's congruence. *Acta Arith.*, 128(4):303–318, 2007.

[SP07] Ling-Ling Shi and Hao Pan. A q-analogue of Wolstenholme's harmonic series congruence. *Amer. Math. Monthly*, 114(6):529–531, 2007.

[Sta97] Richard P. Stanley. *Enumerative Combinatorics, Volume 1*. Cambridge University Press, 1997.

[Wol62] Joseph Wolstenholme. On certain properties of prime numbers. *Quart. J. Math., Oxford Ser.*, 5:35–39, 1862.